

Spectral estimation for Hamiltonians: a comparison between classical imaginary-time evolution and quantum real-time evolution

ME Stroeks[1,2], J Helsen[3] and BM Terhal[1,2,4]

[1] QuTech, Delft University of Technology, P.O. Box 5046, 2600 GA Delft, The Netherlands
[2] Faculty of EEMCS, Delft University of Technology, Mekelweg 4, 2628 CD Delft, The Netherlands
[3] CWI & QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands
[4] JARA Institute for Quantum Information, Forschungszentrum Juelich, D-52425 Juelich, Germany

E-mail: m.e.h.m.stroeks@tudelft.nl, Jonas.Helsen@cwi.nl, b.m.terhal@tudelft.nl

arXiv:2204.01113v2 [quant-ph] 9 May 2022

Abstract. We present a classical Monte Carlo (MC) scheme which efficiently estimates an imaginary-time, decaying signal for stoquastic (i.e. sign-problem-free) local Hamiltonians. The decay rates in this signal correspond to Hamiltonian eigenvalues (with associated eigenstates present in an input state) and can be classically extracted using a classical signal processing method like ESPRIT. We compare the efficiency of this MC scheme to its quantum counterpart in which one extracts eigenvalues of a general local Hamiltonian from a real-time, oscillatory signal obtained through quantum phase estimation circuits, again using the ESPRIT method. We prove that the ESPRIT method can resolve $S = \text{poly}(n)$ eigenvalues, assuming a $1/\text{poly}(n)$ gap between them, with $\text{poly}(n)$ quantum and classical effort through the quantum phase estimation circuits, assuming efficient preparation of the input state. We prove that our Monte Carlo scheme plus the ESPRIT method can resolve $S = O(1)$ eigenvalues, assuming a $1/\text{poly}(n)$ gap between them, with $\text{poly}(n)$ purely classical effort for stoquastic Hamiltonians, requiring some access structure to the input state. However, we also show that under these assumptions, i.e. $S = O(1)$ eigenvalues, assuming a $1/\text{poly}(n)$ gap between them and some access structure to the input state, one can achieve this with $\text{poly}(n)$ purely classical effort for *general* local Hamiltonians. These results thus quantify some opportunities and limitations of classical Monte Carlo methods for spectral estimation of Hamiltonians. We numerically compare the MC eigenvalue estimation scheme (for stoquastic Hamiltonians) and the QPE eigenvalue estimation scheme by implementing them for an archetypal stoquastic Hamiltonian system: the transverse field Ising chain.

Keywords: quantum information, stoquastic Hamiltonian, spectral estimation, Monte Carlo algorithm, quantum phase estimation, ESPRIT.