


SciPost Physics Submission

Holographic Quantum Scars

Diego Liska [1*], Vladimir Gritsev [1,2], Ward Vleeshouwers [1,3], Jiří Minář [1,2,4]

1 Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands
2 QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands
3 QuiX Quantum B.V., Hengelosestraat 500, 7521 AN Enschede, The Netherlands
4 CWI, Science Park 123, 1098 XG Amsterdam, the Netherlands

*d.liska@uva.nl, j.minar@uva.nl

July 6, 2023

arXiv:2212.05962v3 [hep-th] 5 Jul 2023

Abstract

We discuss a construction of quantum many-body scars in the context of holography. We consider two-dimensional conformal field theories and use their dynamical symmetries, naturally realized through the Virasoro algebra, to construct scarred states. By studying their Loschmidt amplitude, we evaluate the states' periodic properties. A geometrical interpretation allows us to compute the expectation value of the stress tensor and entanglement entropy of these scarred states. We show that their holographic dual is related by a diffeomorphism to empty AdS, even for energies above the black hole threshold. We also demonstrate that expectation values in the scarred states are generally non-thermal and that their entanglement entropy grows with the energy as $\log(E)$ in contrast to $\sqrt{E}$ for the typical (bulk) states. Furthermore, we identify fixed points on the CFT plane associated with divergent or vanishing entanglement entropy in the limit where the scarred states have infinite energy.

Contents

1 Introduction **2**
1.1 Quantum many-body scars 2
1.2 Algebraic approach to weakly-nonergodic behavior 3
1.2.1 Dynamical symmetries and scars 3
1.2.2 Quasi-exactly solvable models 4
1.2.3 Floquet scars 4
1.3 Formulation of the problem and relation to other works 5

2 Scars in 2d CFTs **7**
2.1 Loschmidt amplitude 9

3 Scars as reparameterizations **11**
3.1 The stress tensor 13
3.2 Thermalization 13

4 Entanglement **14**
4.1 Scarred states 17