

2022–
2024

Upper Bounds on the Distillable Randomness of Bipartite Quantum States

Ludovico Lami*†, Bartosz Regula‡, Xin Wang§, and Mark M. Wilde¶
* Institute for Theoretical Physics, Korteweg–de Vries Institute for Mathematics, and QuSoft,
University of Amsterdam, the Netherlands. Email: ludovico.lami@gmail.com
† Institute for Theoretical Physics and IQST, University of Ulm, Albert-Einstein-Allee 11, D-89069 Ulm, Germany
‡Department of Physics, Graduate School of Science, The University of Tokyo,
Bunkyo-ku, Tokyo 113-0033, Japan. Email: bartosz.regula@gmail.com
§Institute for Quantum Computing, Baidu Research, Beijing 100193, China. Email: wangxinfelix@gmail.com
¶School of Electrical and Computer Engineering, Cornell University,
Ithaca, New York 14850, USA. Email: wilde@cornell.edu

arXiv:2212.09073v1 [quant-ph] 18 Dec 2022

***Abstract*—The distillable randomness of a bipartite quantum state is an information-theoretic quantity equal to the largest net rate at which shared randomness can be distilled from the state by means of local operations and classical communication. This quantity has been widely used as a measure of classical correlations, and one version of it is equal to the regularized Holevo information of the ensemble that results from measuring one share of the state. However, due to the regularization, the distillable randomness is difficult to compute in general. To address this problem, we define measures of classical correlations and prove a number of their properties, most importantly that they serve as upper bounds on the distillable randomness of an arbitrary bipartite state. We then further bound these measures from above by some that are efficiently computable by means of semi-definite programming, we evaluate one of them for the example of an isotropic state, and we remark on the relation to quantities previously proposed in the literature.**

I. Introduction

The distillable randomness of a bipartite quantum state is equal to the largest net rate at which uniformly random, perfectly correlated bits can be distilled from the state by means of local operations and one-way classical communication (1W-LOCC) [1]. That is, in this scenario, the net rate is the rate at which shared randomness is distilled minus the rate at which classical bits are communicated to accomplish the task. It is important to subtract the classical communication rate, as failure to do so would trivialize the task, allowing for an infinite number of random bits to be shared. This task fundamentally has its roots in classical information theory [2], [3]. Here we also extend the task to allow for general local operations and classical communication (LOCC).

The distillable randomness of a bipartite state is considered a fundamental measure of classical correlations contained in a state. There has been a large literature on this topic, starting with [4], [5] and reviewed in [6], due to the wide interest in understanding correlations present in quantum states. However, [1] is the one of the few papers to consider understanding these correlations in an information-theoretic manner (see also [7]–[13] for other perspectives and related works).

Ref. [1] provided a formal solution for the aforementioned information-theoretic task. However, like many such formal solutions in quantum Shannon theory, it does not provide a computationally efficient procedure for quantifying correlations because it is expressed as a multi-letter or regularized formula. Our goal here is to fill this void by providing upper bounds on the distillable randomness that are efficiently computable by semi-definite programming. We also justify how some of the newly proposed measures are themselves measures of classical correlations contained in a bipartite state.

Although our results here can be viewed as a static counterpart to the results in the dynamical setting from [14]–[17], their derivation requires some new insights, as we explain in detail below. Ref. [17] proved that generalizations of the dynamical measures from [14]–[16] give upper bounds on the rate at which classical messages be communicated from one party to another, whenever they have access to a bipartite channel as a communication resource. A special case of this setting is when a sender is communicating classical messages to a receiver, with the assistance of a classical feedback channel. In contrast, as described earlier, our static setting here involves two parties distilling shared randomness from a bipartite state, by means of 1W-LOCC or general LOCC.

We comment here on the novel contributions of our paper when compared to earlier works such as [14]–[17]. The main measures of classical correlations proposed here, denoted by γ, C_γ, and $\mathbf{\Gamma}$ in Section III, are inspired by the measures that have appeared earlier in [14]–[17], denoted there by β, C_β, and $\boldsymbol{\Upsilon}$. However, the measures proposed here are symmetric with respect to a swap of the A and B systems. This property is critical for bounding the LOCC-assisted distillable randomness from above, as accomplished in Theorems 11 and 12. Furthermore, we prove here that these measures obey the classical communication bound in Proposition 7, also known as a "non-lockability" property in prior work on correlation measures [18], [19]. Finally, Lemma 9 and Proposition 10 are critical bounds that allow us to compare the actual state at the end of a randomness distillation protocol with an ideal state, and they ultimately lead to a strong converse upper bound for the distillable randomness. For the task of randomness distillation, the target (ideal) state is a mixed state, which is in