arXiv:2302.11499v2 [quant-ph] 1 Nov 2023

Teleportation of quantum coherence

Sohail [1], Arun K Pati [2], Vijeth Aradhya [3,4], Indranil Chakrabarty [2,3], Subhasree Patro [3,5,61]

[11] *Quantum Information and Computation Group, Harish-Chandra Research Institute, A CI of Homi Bhabha National Institute, Chhatnag Road, Jhunsi, Prayagraj - 211019, India.*
[2] *Centre for Quantum Science and Technology, International Institute of Information Technology-Hyderabad, Gachibowli, Hyderabad, India.*
[3] *Center for Security, Theory and Algorithmic Research International Institute of Information Technology-Hyderabad, Gachibowli, Hyderabad, India.*
[4] *Department of Computer Science, National University of Singapore - 117417 Singapore.*
[5] *Utrecht University, 3584 CS Utrecht, The Netherlands*
[6] *QuSoft, Centrum Wiskunde & Informatica, 1098 XG Amsterdam, The Netherlands.*

We investigate whether it is possible to teleport the coherence of an unknown quantum state from Alice to Bob by communicating a lesser number of classical bits in comparison to what is required for teleporting an unknown quantum state. We find that we cannot achieve perfect teleportation of coherence with one bit of classical communication for an arbitrary qubit. However, we find that if the qubit is partially known, i.e., chosen from the equatorial and polar circles of the Bloch sphere, then teleportation of coherence is possible with the transfer of one cbit of information when we have maximally entangled states as a shared resource. In the case of the resource being a non-maximally entangled state, we can teleport the coherence with a certain probability of success. In a general teleportation protocol for coherence, we derive a compact formula for the final state at Bob's lab in terms of the composition of the completely positive maps corresponding to the shared resource state and joint POVM performed by Alice on her qubit and the unknown state. Using this formula, we show that teleportation of the coherence of a partially known state with real matrix elements is possible perfectly with the help of a maximally entangled state as a resource. Furthermore, we explore the teleportation of coherence with the Werner states and show that even when the Werner states become separable, the amount of teleported coherence is non-zero, implying the possibility of teleportation of coherence without entanglement.

I. INTRODUCTION

Quantum Coherence [1] and entanglement [2] are two central features in quantum theory that make the theory counterintuitive and can also be utilised as a resource to perform important information theoretic and computational tasks. Motivated by this increasing importance, a general study from the perspective of resource theory is being formulated. Several measures based on resource theoretic approaches are available to quantify the coherence and the entanglement present in a quantum system [1–19]. Quantum coherence can be interpreted in several ways, like as a measure of non-classicality in physical systems, a measure of superposition, and as a quantity capturing the wave aspect of the state vector [20–22]. It is also observed that perfect cloning and broadcasting of quantum coherence are not possible [23, 24]. Coherence acts as a resource in tasks like quantum algorithms [25–27], biological processes [28, 29] quantum metrology [30–34], reference frame alignment [35], thermodynamic tasks [36–40]. From the point of view of resource theory, coherence is classified into two classes, namely speakable coherence and unspeakable coherence[41]. The quantification of coherence involves a chosen basis. In situations where the labelling of the chosen basis is not important, the relevant notion of coherence is speakable one, and in situations where the labelling or the identity of the basis elements matters, the relevant notion of coherence is unspeakable one. Coherence involved in quantum metrology, reference frame alignment, and thermodynamic tasks are examples of unspeakable coherence, whereas the coherence involved in computational, cryptographic, and communication tasks are examples of speakable coherence.

In the last two decades, several developments have been made in quantum based technologies. These include quantum repeaters for communicating over large distances [42], quantum teleportation [43], broadcasting of entanglement [44–47]. These were introduced in the process with the vision of having quantum networks [48–50]. In particular, "Quantum teleportation" is one of the most fascinating discoveries of the 20th century [43]. The entangled states which are useful for teleportation are identified as well as detected in this process [51–54]. Research was also done to implement the teleportation process, if not perfectly but at least with a certain probability of success [55]. Recently, it has also been utilized to understand phenomena like closed time-like curves [56]. It has been demonstrated in the laboratory with the help of various resources like photonic qubits, nuclear magnetic resonance (NMR)[57], optical modes [58–66] , atomic ensembles [67–70], trapped atoms [71–75], and solid state systems[76–79]. Efficient teleportation was also achieved in terms of distance [80, 81].

In this article, in general, we address the question of whether we can teleport the coherence of a quantum state, with a lesser number of classical bits than what is required to teleport the state itself. It is already known that teleportation of an unknown state is possible with the transfer of two classical bits of information. The process of teleportation of a known state is known as remote state preparation, and it requires the transfer of one bit of information for the process to be (only for equatorial qubit) perfect [82]. It is obvious that once the entire information of the state is recreated in a different location, the coherence of the state is also transferred. There are many instances where it has been seen that the coherence