arXiv:2305.16803v4 [quant-ph] 30 Jan 2025

Quantum work extraction efficiency for noisy quantum batteries: the role of coherence

Salvatore Tirone,[1,2,3,*] Raffaele Salvia,[1,†] Stefano Chessa,[4,5,‡] and Vittorio Giovannetti[4]

[1] *Scuola Normale Superiore, I-56126 Pisa, Italy*
[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[3] *Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*
[4] *NEST, Scuola Normale Superiore and Istituto Nanoscienze-CNR, I-56126 Pisa, Italy*
[5] *Electrical and Computer Engineering, University of Illinois Urbana-Champaign, Urbana, Illinois, 61801, USA*

Quantum work capacitances and maximal asymptotic work/energy ratios are figures of merit characterizing the robustness against noise of work extraction processes in quantum batteries formed by collections of quantum systems. In this paper we establish a direct connection between these functionals and, exploiting this result, we analyze different types of noise models mimicking self-discharging (amplitude damping channels), thermalization (generalized amplitude damping channels) and dephasing effects (simultaneous phase-amplitude damping channels). In this context we show that input quantum coherence can significantly improve the storage performance of noisy quantum batteries and that the maximum output ergotropy is not always achieved by the maximum available input energy.

I. INTRODUCTION

The use of quantum effects have been proposed as useful methods to speed up charging processes of batteries [1–9]. To assess the stability of these proposals it is crucial to analyze how such schemes perform in the presence of environmental noise, as this represents a more realistic scenario, as it has been proved by the first experimental implementations [10–15], and can have a significant impact on the efficiency of energy recovery. For noiseless models capacities for Quantum Batteries (QBs) have been proposed [7, 16, 17] while, from a resource theoretical point of view, the thermodynamic capacity (in the sense of process simulability) of quantum channels has been defined [18]. In the presence of noise, in terms of energetic manipulation, a few results have been obtained in specific frameworks, see e.g. [19–35]. In an effort to generalize and facilitate the comparison between different physical platforms, two types of universal figures of merit, i.e. the quantum work capacitances and the maximal asymptotic work/energy ratios (MAWERs), have been introduced in [36].These quantities gauge different aspects of the efficiency of work extraction from noisy quantum battery systems providing a parallel point of view over the concept of efficiency of other quantum thermodynamic objects [37–39], such as e.g. quantum heat engines, which model alternative thermodynamic operational protocols [40–42]. Here, the aforementioned new figures of merit are useful in scenarios where one has at disposal large collections of identical energy storing quantum devices: quantum cells or q-cells in brief. Specifically, quantum work capacitances target the maximum work that can be recovered per q-cell assuming that on average each cell allocates a fixed portion of the total energy E initially stored in the overall battery. For MAWERs, on the contrary, the work extraction efficiency is gauged treating the q-cells as a free resource, assuming their number n to be much larger than the minimal amount that could host the full input energy E. The evaluations of quantum work capacitances and of MAWERs rely on complex and resource-constrained optimization problems similar to those one faces in quantum communication [43–49], where encoding/decoding strategies must be designed to mitigate the effects of dissipation and decoherence – see Fig. 1. Closed formulas that enable one to explicitly compute the value of some of the work capacitances have been recently presented in Ref. [50]. Here we extend these findings to the MAWERs, proving that the latter can be expressed as a special limit of the associated quantum work capacitance functionals. Using these results we next investigate several models of noisy batteries describing a wide range of deterioration effects that are physically relevant, including self-discharging, thermalization, and dephasing. In this context we find that the work that can be extracted from the QB is not always a monotonic function of the energy that was initially stored in the system. As a consequence we show that, for some noise models, an incompletely charged battery may perform better than a fully charged one. Additionally, we observe that the presence of quantum coherence strengthens the resistance of QBs against environmental noise. Specifically, when it comes to self-discharging and thermal noise, we show that coherent input states outperform all conceivable classical (i.e. incoherent) initial states of the battery. This result strengthens the evidences built up in the literature in recent years, both in the noisy and the noiseless settings [51–55], that quantum coherence can provide an advantage in the quantum setting for energy manipulation tasks and proves to be a useful resource as happens for information processing applications [56–59].

* s.tirone@uva.nl
† raffaele.salvia@sns.it
‡ schessa@illinois.edu