

2022–
2024

Entanglement cost for infinite-dimensional physical systems

Hayata Yamasaki,[1,*] Kohdai Kuroiwa,[2,3,†] Patrick Hayden,[4,‡] and Ludovico Lami[5,6,7,§]

[1]*Department of Physics, Graduate School of Science, The Univerisity of Tokyo, 7–3–1 Hongo, Bunkyo-ku, Tokyo, 113–0033, Japan*
[2]*Institute for Quantum Computing and Department of Combinatorics and Optimization, University of Waterloo, Ontario, Canada, N2L 3G1*
[3]*Perimeter Institute for Theoretical Physics, Ontario, Canada, N2L 2Y5*
[4]*Stanford Institute for Theoretical Physics, Stanford University, Stanford, California 94305, USA*
[5]*QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*
[6]*Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105–107, 1098 XG Amsterdam, The Netherlands*
[7]*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

arXiv:2401.09554v1 [quant-ph] 17 Jan 2024

We prove that the entanglement cost equals the regularized entanglement of formation for any infinite-dimensional quantum state ρ_{AB} with finite quantum entropy on at least one of the subsystems A or B. This generalizes a foundational result in quantum information theory that was previously formulated only for operations and states on finite-dimensional systems. The extension to infinite dimensions is nontrivial because the conventional tools for establishing both the direct and converse bounds, i.e., strong typically, monotonicity, and asymptotic continuity, are no longer directly applicable. To address this problem, we construct a new entanglement dilution protocol for infinite-dimensional states implementable by local operations and a finite amount of one-way classical communication (one-way LOCC), using weak and strong typicality multiple times. We also prove the optimality of this protocol among all protocols even under infinite-dimensional separable operations by developing an argument based on alternative forms of monotonicity and asymptotic continuity of the entanglement of formation for infinite-dimensional states. Along the way, we derive a new integral representation for the quantum entropy of infinite-dimensional states, which we believe to be of independent interest. Our results allow us to fully characterize an important operational entanglement measure — the entanglement cost — for all infinite-dimensional physical systems.

CONTENTS

I. Introduction: the asymptotic continuity catastrophe 2
A. Entanglement dilution 2
B. Enter infinite-dimensional systems 3
C. Main contribution 4

II. Preliminaries 5
A. Quantum mechanics 5
B. Topologies on the set of quantum channels 6
C. LOCC in infinite dimensions 7
D. Entanglement cost 9
E. Entanglement of formation 11
F. What exactly goes wrong in infinite dimensions? 13

* hayata.yamasaki@gmail.com
† kkuroiwa@uwaterloo.ca
‡ phayden@stanford.edu
§ ludovico.lami@gmail.com