

2022–
2024

Quantum communication on the bosonic loss-dephasing channel

Francesco Anna Mele,[1,*] Farzin Salek,[2,†] Vittorio Giovannetti,[1,‡] and Ludovico Lami[3,4,5,§]

[1]*NEST, Scuola Normale Superiore and Istituto Nanoscienze, Consiglio Nazionale delle Ricerche, Piazza dei Cavalieri 7, IT-56126 Pisa, Italy*
[2]*Department of Mathematics, Technical University of Munich, Boltzmannstrasse 3, 85748 Garching, Germany*
[3]*QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[4]*Korteweg-de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*
[5]*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*

Quantum optical systems are typically affected by two types of noise: photon loss and dephasing. Despite extensive research on each noise process individually, a comprehensive understanding of their combined effect is still lacking. A crucial problem lies in determining the values of loss and dephasing for which the resulting loss-dephasing channel is anti-degradable, implying the absence of codes capable of correcting its effect or, alternatively, capable of enabling quantum communication. A conjecture in [Quantum 6, 821 (2022)] suggested that the bosonic loss-dephasing channel is not anti-degradable if the loss is below 50%. In this paper we refute this conjecture, specifically proving that for any value of the loss, if the dephasing is above a critical value, then the bosonic loss-dephasing channel is anti-degradable. While our result identifies a large parameter region where quantum communication is not possible, we also prove that if two-way classical communication is available, then quantum communication — and thus quantum key distribution — is always achievable, even for high values of loss and dephasing.

arXiv:2401.15634v3 [quant-ph] 25 Jul 2024

I. INTRODUCTION

Quantum optical platforms are key elements of quantum technologies, contributing significantly to both quantum communication and quantum computation [1–9]. Since the potential benefits of quantum technologies are hindered by the presence of decoherence [10], the investigation of decoherence sources affecting bosonic systems and the development of bosonic quantum error-correcting codes have been extensively analysed in recent years [11–22]. The primary noise processes in bosonic systems that act as dominant sources of decoherence are *photon loss* and *bosonic dephasing* [23–25], which have been both extensively analysed [13, 26–28]. Loss affects the system by causing it to dissipate some of its energy, whereas dephasing works to transform coherent superpositions into probabilistic mixtures. Although loss and dephasing sources can simultaneously affect bosonic systems [15, 29], such as in superconducting systems [30, 31], the existing literature provides only partial results about their combined effect [32]. On a technical level, understanding the combined effect of loss and dephasing is challenging due to the conflicting behaviours they exhibit: the action of loss takes a simple form when written in the coherent state basis but is complicated to analyse in the Fock basis [2], whereas dephasing demonstrates the opposite pattern, making the analysis of their combined effect quite intricate.

Consider an optical link (e.g. an optical fibre or a free-space link) or a quantum memory affected by both loss and dephasing, where the link is used for quantum communication and the memory for quantum computation. A crucial challenge is to determine the conditions under which there exist protocols capable of enabling reliable quantum communication across the optical link or capable of mitigating the combined noise affecting the quantum memory. This problem is closely related to the anti-degradability condition in quantum Shannon theory: if a noise channel is *anti-degradable* [33, 34], there are no quantum communication protocols for reliable information transmission or quantum error-correcting codes capable of overcoming it. Consequently, it is crucial to understand whether the combined effect of loss and dephasing results in an anti-degradable channel. This has been a puzzling problem, to the point that in [32] it was conjectured that the combined loss-dephasing noise does not result in an anti-degradable channel if the loss is below 50%.

In this paper we refute the above conjecture; specifically, we prove that for any value of the photon loss there exists a *critical value* of the dephasing above which the resulting *bosonic loss-dephasing channel* is anti-degradable. Our discovery thus identifies a large region of the loss-dephasing parameter space where correcting the noise and achieving reliable quantum communication is impossible. On the more positive side, however, we also prove that if the sender and the receiver are assisted by two-way classical communication, then reliable quantum communication — and thus quantum key distribution — is always possible, even in scenarios characterised by arbitrarily high levels of loss and dephasing. Ours are the first analytical results to characterise the transmission of quantum information in the presence of both loss and dephasing noise.

The structure of the paper is as follows. In Section II, we briefly review some preliminary notions necessary for stating our results. In Section III, we present our results on the anti-degradability of the bosonic