

2022–
2024

# Learning quantum states of continuous-variable systems


Francesco A. Mele,[1,*] Antonio A. Mele,[2,†] Lennart Bittel,[2,†] Jens Eisert,[2,3,‡] Vittorio Giovannetti,[4,*] Ludovico Lami,[5,6,7,‡] Lorenzo Leone,[2,†] and Salvatore F. E. Oliviero[4,*]

[1]*NEST, Scuola Normale Superiore and Istituto Nanoscienze, Piazza dei Cavalieri 7, IT-56126 Pisa, Italy*
[2]*Dahlem Center for Complex Quantum Systems, Freie Universität Berlin, 14195 Berlin, Germany*
[3]*Helmholtz-Zentrum Berlin für Materialien und Energie, Berlin, Germany*
[4]*NEST, Scuola Normale Superiore and Istituto Nanoscienze, Consiglio Nazionale delle Ricerche, Piazza dei Cavalieri 7, IT-56126 Pisa, Italy*
[5]*QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[6]*Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*
[7]*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*


arXiv:2405.01431v4 [quant-ph] 7 Nov 2024


Quantum state tomography, aimed at deriving a classical description of an unknown state from measurement data, is a fundamental task in quantum physics. In this work, we analyse the ultimate achievable performance of tomography of continuous-variable systems, such as bosonic and quantum optical systems. We prove that tomography of these systems is extremely inefficient in terms of time resources, much more so than tomography of finite-dimensional systems: not only does the minimum number of state copies needed for tomography scale exponentially with the number of modes, but it also exhibits a dramatic scaling with the trace-distance error, even for low-energy states, in stark contrast with the finite-dimensional case. On a more positive note, we prove that tomography of Gaussian states is efficient. To accomplish this, we answer a fundamental question for the field of continuous-variable quantum information: if we know with a certain error the first and second moments of an unknown Gaussian state, what is the resulting trace-distance error that we make on the state? Lastly, we demonstrate that tomography of non-Gaussian states prepared through Gaussian unitaries and a few local non-Gaussian evolutions is efficient and experimentally feasible.


## I. INTRODUCTION

A fundamental task in quantum physics, known as *quantum state tomography* [1], is to derive a classical description of a quantum system from experimental data. It serves not only as a powerful method to investigate Nature, but also as a diagnostic tool for benchmarking and verifying quantum devices [1–4]. The concept of tomography dates back to the 1990s [5, 6], when it was first introduced for *continuous-variable* (CV) systems [7] and then for finite-dimensional systems. Over the years, tomography algorithms for CV systems have been extensively developed, becoming a bread-and-butter tool for quantum opticians [5, 6].

In the last decade, rapid advances in accurately preparing quantum states [8] have spurred the development of a new research field known as *quantum learning theory* [1], which investigates the question of how to learn properties of quantum systems as efficiently as possible. When the focus is on obtaining a full classical description of a quantum system, such investigation reduces to quantum state tomography, which is considered the cornerstone of quantum learning theory [1]. Despite tomography being a longstanding concept, many of its fundamental properties have only recently been unveiled due to advancements in this new field, albeit only for finite-dimensional systems and not for CV systems.

Let us introduce the concept of quantum state tomography from a quantum learning theory perspective. Formally, the task of quantum state tomography involves two parameters: the number of state copies $N$ and the error parameter $\varepsilon$. Given $N$ copies of an unknown state, the goal is to output a classical description of a state which is guaranteed to be $\varepsilon$-close to the true unknown state with high probability (this is stated in all rigour in the Methods section). Among various notion of distance to measure the $\varepsilon$-closeness, the *trace distance* emerges as the most meaningful one due to its operational significance [9, 10]: given two states $\varepsilon$-close in trace distance, no quantum measurement can distinguish them by more than $\varepsilon$. Thus, the two states are in fact indistinguishable to any physical observer within a $\varepsilon$ resolution. The optimal performance of tomography is quantified by the *sample complexity*, which is the minimum number of copies $N$ needed to achieve tomography with error $\varepsilon$. When the sample complexity and the running time of the tomography procedure scale polynomially with the number of constituents (e.g., qubits, modes), tomography is deemed *efficient*.

With vast many-body systems naturally emerging in Nature on one side, and the rapid advancement in constructing large-scale quantum devices on the other, minimising required resources has become imperative. Consequently, determining the sample complexity of tomography of physically relevant classes of states is currently a pressing problem [1]. Notably, the sample complexity has been determined for $n$-qubit systems [1, 11–14]: tomography of mixed states requires $O(4^n/\varepsilon^2)$ copies, while for pure states the required number of copies reduces to $O(2^n/\varepsilon^2)$.

While quantum learning theory has been extensively developed for finite-dimensional systems, it remains almost unex-


* {francesco.mele, vittorio.giovannetti, salvatore.oliviero}@sns.it
† {a.mele, l.bittel, lorenzo.leone}@fu-berlin.de
‡ {jenseisert, ludovico.lami}@gmail.com