
2021–
2026


2021–
2025



arXiv:2405.03517v1 [math.CO] 6 May 2024

All S_p notions of quantum expansion are equivalent

Francisco Escudero Gutiérrez* Garazi Muguruza†

Abstract

In a recent work Li, Qiao, Wigderson, Wigderson and Zhang introduced notions of quantum expansion based on S_p norms and posed as an open question if they were all equivalent. We give an affirmative answer to this question.

1 Introduction

In this note we consider the problem of showing that certain notions of quantum expansion are equivalent. Quantum expansion is an analogue of the well-studied concept of graph expansion. The latter has many equivalent versions that let it play an important role in many fields such as geometry, group-theory, combinatorics and theoretical computer science (for a survey see [HLW06]).

Given a d-regular graph $G = ([n], E)$, its *edge expansion* is given by

$$h(G) = \min_{W \subset [n], |W| \leq n/2} \frac{|\partial W|}{d|W|},$$

where $|W|$ stands for the size of W and $|\partial W|$ is the number of edges that go from W to its complement. Let $\mathbf{B} = (B_1, \ldots, B_d)$ be permutation matrices that decompose the adjacency matrix A of G as $\sum_{i\in[n]} B_i$[1]. In particular, $\mathbf{B}$ is a bistochastic tuple, meaning that $\sum_{i\in[n]} B_i^* B_i = \sum_{i\in[n]} B_i B_i^* = d \cdot \mathrm{Id}$. Thus, it induces a quantum unital channel via

$$\Phi_{\mathbf{B}}(X) = \frac{1}{d} \sum_{i\in[n]} B_i X B_i^*.$$

A quantum channel is a linear operator from $M_n(\mathbb{C})$ to $M_n(\mathbb{C})$ that maps quantum states (positive semidefinite matrices with trace 1) to quantum states. It is unital if it maps the identity to itself. Note that if $\mathbf{B}$ is a permutation tuple, W a subset of $[n]$ and P_W the projector onto the space spanned by the canonical basis vectors indicated by W, then $\langle \mathrm{Id} - P_W, \Phi_{\mathbf{B}}(P_W)\rangle = |\partial W|$. Here $\langle \cdot, \cdot \rangle$ stands for the inner product $\langle M, N\rangle = \mathrm{Tr}[M^* N]$. Motivated by this, Hastings introduced the *quantum edge expansion* of a bistochastic tuple $\mathbf{B} \in M_n(\mathbb{C})^d$, which is given by

$$h_Q(\mathbf{B}) = \min_{V < \mathbb{C}^n, \dim V \leq n/2} \frac{\langle \mathrm{Id} - P_V, \Phi_{\mathbf{B}}(P_V)\rangle}{d \dim V} \tag{1}$$

*Qusoft and CWI feg@cwi.nl
†Qusoft and UvA g.muguruzalasa@uva.nl

[1]It follows from Hall's theorem that this decomposition always exists.