

2022–
2024

Computable entanglement cost under positive partial transpose operations

Ludovico Lami,[1,2,3,4,*] Francesco Anna Mele,[5,†] and Bartosz Regula[6,‡]

[1]*Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy*
[2]*QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[3]*Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*
[4]*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[5]*NEST, Scuola Normale Superiore and Istituto Nanoscienze, Consiglio Nazionale delle Ricerche, Piazza dei Cavalieri 7, IT-56126 Pisa, Italy*
[6]*Mathematical Quantum Information RIKEN Hakubi Research Team, RIKEN Cluster for Pioneering Research (CPR) and RIKEN Center for Quantum Computing (RQC), Wako, Saitama 351-0198, Japan*

arXiv:2405.09613v2 [quant-ph] 7 Mar 2025

Quantum information theory is plagued by the problem of regularisations, which require the evaluation of formidable asymptotic quantities. This makes it computationally intractable to gain a precise quantitative understanding of the ultimate efficiency of key operational tasks such as entanglement manipulation. Here we consider the problem of computing the asymptotic entanglement cost of preparing noisy quantum states under quantum operations with positive partial transpose (PPT). By means of an analytical example, a previously claimed solution to this problem is shown to be incorrect. Building on a previous characterisation of the PPT entanglement cost in terms of a regularised formula, we construct instead a hierarchy of semi-definite programs that bypasses the issue of regularisation altogether, and converges to the true asymptotic value of the entanglement cost. Our main result establishes that this convergence happens exponentially fast, thus yielding an efficient algorithm that approximates the cost up to an additive error ε in time $\mathrm{poly}\big(D, \log(1/\varepsilon)\big)$, where D is the underlying Hilbert space dimension. To our knowledge, this is the first time that an asymptotic entanglement measure is shown to be efficiently computable despite no closed-form formula being available.

Introduction.— Quantum Shannon theory studies the fundamental limitations on the manipulation of quantum information in the presence of external noise. Calculating those limits often involves computing certain functions that encapsulate the ultimate capabilities of information carriers. Paradigmatic examples include the various capacities of quantum channels, such as the classical [1, 2], quantum [3–5], private [5, 6], and entanglement-assisted [7, 8] capacities, but also the operational entanglement measures that tell us how much entanglement can be extracted from a given bipartite quantum state, i.e. the distillable entanglement [9–13], and vice versa, how much entanglement must be invested to *create* that state [14–17]. This latter quantity, the entanglement cost, is the main focus of this work.

With the sole exception of the entanglement-assisted capacity, all of the above functions are expressed by *regularised formulas*, i.e. formulas that involve an explicit limit $n \to \infty$ over the number of uses of the channel or the available copies of the state. For example, by the Lloyd–Shor–Devetak theorem [3–5] the quantum capacity of a channel $\mathcal{N}$ equals $Q(\mathcal{N}) = \lim_{n\to\infty} \frac{1}{n} I_c\big(\mathcal{N}^{\otimes n}\big)$, where $I_c(\mathcal{N})$ is the 'coherent information' of $\mathcal{N}$, and $\mathcal{N}^{\otimes n}$ represents n parallel uses of $\mathcal{N}$. In stark contrast with classical information theory, for quantum channels it holds in general that $I_c(\mathcal{N}^{\otimes n}) \neq n I_c(\mathcal{N})$, meaning that evaluating the limit cannot be avoided. Such non-additivity is a fundamental feature of most settings encountered in quantum information [18–23]. Analogously, the entanglement cost of preparing a state $\rho = \rho_{AB}$ using local operations and classical communication (LOCC) is given by

$$E_{c,\mathrm{LOCC}}(\rho) = \lim_{n\to\infty} \frac{1}{n} E_f\big(\rho^{\otimes n}\big)\,, \tag{1}$$

where E_f is the 'entanglement of formation' [11, 24]. The precise nature of these formulas is not so important here; what is important, however, is that the regularisation $n \to \infty$ makes them analytically hard to control and computationally intractable. Indeed, on the one hand the dimension of the quantum system on which $\mathcal{N}^{\otimes n}$ acts, or to which $\rho^{\otimes n}$ pertains, is exponential in n, quickly rendering numerical calculations infeasible as n grows; on the other, there is no guarantee on the quality of the approximation obtained by stopping at the nth level in any of these formulas — for instance, an unbounded n may be required to even check that the quantum capacity is non-zero [25]. The regularisation thus appears to be an omnipresent curse that stifles almost every attempt to quantitatively understand the ultimate limitations of quantum information manipulation.

But is that really so? In this work we show how to overcome this fundamental obstacle in a specific case, by efficiently calculating a type of entanglement cost — expressed as a regularised quantity — on all quantum states. To this end, we look at a problem that has been studied by many authors [26–30], but for which a full solution had not been found prior to our work: namely, zero-error asymptotic entanglement cost under 'positive partial transpose' (PPT) operations. The roots of PPT transformations lie in the fundamental connection between entanglement and partial transposition identified by Peres [31] and the Horodeckis [32, 33]. In the context of entanglement manipulation, PPT operations were introduced in the pioneering works of Rains [34, 35] as a mathematically natural relaxation of the much more complicated set of LOCC operations, providing a convenient way to gain some insights into the latter and, importantly, giving hope for an easier understanding of