

Multilevel Electromagnetically Induced Transparency Cooling

Katya Fouka[1,2],* Athreya Shankar[3,4], Ting Rei Tan[5,6,7], and Arghavan Safavi-Naini[1,2]

[1]*Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*
[2]*QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*
[3]*Department of Physics, Indian Institute of Technology Madras, Chennai 600036, India*
[4]*Center for Quantum Information, Communication and Computing, Indian Institute of Technology Madras, Chennai 600036, India*
[5]*School of Physics, University of Sydney, NSW 2006 Sydney, Australia*
[6]*ARC Centre of Excellence for Engineered Quantum Systems, University of Sydney, NSW 2006, Australia*
[7]*Sydney Nano Institute, University of Sydney, NSW 2006, Australia*

(Dated: November 27, 2025)

Electromagnetically Induced Transparency (EIT) cooling is a well-established method for preparing trapped ion systems in their motional ground state. However, isolating a three-level system, as required for EIT cooling, is often challenging or impractical. Nonetheless, multilevel systems can inherently host dark states. In this work, we extend the EIT cooling framework to such multilevel systems. We develop a formalism to accurately determine the cooling rate in the weak sideband coupling regime and provide an approximate estimate for cooling rates beyond this regime, without the need for explicit simulation of the motional degree of freedom. We clarify the connection between the cooling rate and the absorption spectrum, offering a pathway for efficient near-ground-state cooling of ions with complex electronic structures.

arXiv:2506.14546v2 [physics.atom-ph] 26 Nov 2025

I. INTRODUCTION

Trapped ions have played an important role in advancing quantum simulation, computation, sensing, and metrology [1–3]. These applications typically require the ability to manipulate the motional degrees of freedom of the ions. For example, cooling the ion crystal to its motional ground state is necessary for high gate fidelities [4, 5] as well as enabling quantum enhanced metrology and sensing, which often requires preparation of multi-partite entangled states [6, 7]. In particular, phonon mediated interactions between ions, arising from the coupling of the atomic states with the collective motional degrees of freedom of the ion crystal, are crucial in the operation of logical gates and state preparation protocols [1, 8–10]. However, large ion crystals, required for useful quantum operations, present significant challenges. As the crystal size increases, the corresponding growth of motional modes demands cooling over a broad bandwidth to counteract environmental heating and ensure high-fidelity operations [5, 11, 12].

Experiments with trapped ions typically begin with a Doppler cooling stage [13], followed by sub-Doppler cooling techniques to overcome the temperature limitation imposed by the linewidth of the addressed transition [14]. Commonly used ground-state cooling techniques include Sisyphus cooling [15–18], resolved sideband cooling [19–21] and electromagnetically induced transparency (EIT) cooling [22–24]. Although Sisyphus cooling is capable of simultaneously cooling a broad spectrum of motional modes, its cooling limit is not sufficiently low [15–18]. On the other hand, resolved sideband cooling provides a low cooling limit. However, its narrow cooling bandwidth results in an overall cooling duration that increases with the number of ions [25, 26]. In contrast, EIT cooling offers both fast near ground state preparation and a broad cooling bandwidth, positioning it as one of the most attractive sub-Doppler cooling techniques [27–34].

Although *single* EIT cooling [22–24] is predominantly linked to a closed three-level system, its applicability has been successfully extended to more complex configurations. In particular, EIT cooling has been demonstrated in closed four-level systems, referred to as *double* EIT cooling [32, 33, 35–37]. Additionally, it has been recently performed in multilevel systems by engineering an open three-level system using additional laser sources beyond those required for cooling [38] as well as in an open four-level system [39].

The core principle of EIT cooling relies on using the dark states of the light-atom system. Multilevel systems can naturally support dark states as long as the coherent population trapping condition is met [40]. This suggests that EIT cooling can be performed without the necessity of isolating a three-level system, whether open or closed.

In this paper, we numerically investigate *multilevel* EIT cooling in systems hosting dark states. We begin, in Sec. II., by briefly recalling the principles of single EIT cooling. We summarize established techniques for estimating cooling rates while introducing an alternative method. Furthermore, we assess the validity of these techniques within the sideband coupling regimes. In Sec. III., we model the multilevel cooling transition of a realistic ion for arbitrary nuclear spin I and study multilevel EIT cooling while testing the cooling estimation techniques for $I = 1$ and $I = 7$.

* s.k.fouka@uva.nl