


Ultracold collisions of a neutral atom with a trapped ion in 1D

Seth T. Rittenhouse,[1,2,3] Lorenzo Oghittu,[3,4] Arghavan Safavi-Naini,[3,4] Rene Gerritsma,[5] and Nirav P. Mehta[6,3]

[1] *Department of Physics, the United States Naval Academy, Annapolis, MD, 21402, USA*
[2] *ITAMP, Center for Astrophysics | Harvard & Smithsonian, Cambridge, USA*
[3] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[4] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[5] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, 1098 XH Amsterdam, the Netherlands*
[6] *Department of Physics and Astronomy, Trinity University, San Antonio Texas 78212, USA*

arXiv:2506.15804v1 [physics.atom-ph] 18 Jun 2025

We present a fully quantum mechanical description of a free ^{6}Li atom scattering from a trapped ^{171}Yb$^+$ ion in one dimension. By reformulating the system in polar coordinates and employing the adiabatic representation, we extract a set of coupled adiabatic potentials representing the atom interacting with the ion in different trap states. In an approach similar to quantum defect theory (QDT), we leverage the vast difference in energy scale between the interaction, the trap, and the scattering energy to encapsulate the short-range atom-ion scattering behavior in a single phase parameter. The presence of trapped $(^{171}\text{Yb}^6\text{Li})^+$ molecular-ion states leads to a series of roughly evenly spaced resonances in the scattering cross section. The predicted distribution of resonances at low collision energies is at odds with the expectation of quantum chaos and the Bohigas-Giannoni-Schmit (BGS) conjecture.

I. INTRODUCTION

Hybrid systems consisting of a trapped ion embedded in an ultracold gas of neutral atoms offer exciting prospects in quantum science [1–6]. Such systems benefit from the ultracold temperatures achieved in atomic gases and the ability to individually address the spatially localized trapped ions. Atom-ion interactions serve to entangle the constituent systems, paving the way for several exciting experimental advances, including the observation of atom-ion collisions [7, 8], magnetic Feshbach resonances [9], realizations of sympathetic [1, 10, 11] and swap [12] cooling of the ion by the cold buffer gas, and the study of ultracold atom-ion chemistry [13–15].

A comprehensive understanding of atom-ion hybrid systems hinges on the development of a robust collision theory that is capable of incorporating the confinement of the ion by a radio-frequency (RF) Paul trap. To date, progress has been made considering classical trajectories [16], which revealed that micromotion heating of the ion can be minimized when the atomic mass m_a is chosen to be much smaller than the mass of the ion, m_i. More recently, classical calculations have helped interpret experiments observing trap-assisted formation of long-lived atom-ion complexes [17, 18]. Moreover, classical trajectories exhibit signatures of chaos [19, 20] linked to such states reminiscent of the "sticky collisions" predicted [21–23] and observed [24] in ultracold molecular gases.

While rigorous quantum-mechanical theories considering collisions of atoms and ions in free space have been developed, both at higher thermal energies [25] and in the ultracold regime [26, 27], there is yet to appear a fully quantum model that incorporates the ion confinement. Even at the level of the so-called secular approximation in which ion micromotion is neglected, the harmonic trapping potential significantly complicates the collision physics: the trap center acts as an infinitely massive third particle, and one must consider the system as an effective three-body problem.

As a consequence, new dynamical processes—akin to confinement-induced resonances [28]—arise that are only present when the ion confinement is considered. The most significant of these, and the focus of this work, is the formation of a trapped atom-ion complex during the collision. The binding energy from the formation of the molecular ion propels its center of mass into a high-lying state of the oscillator, and the process is resonantly enhanced if such a state lies near the collision energy. Of particular interest in this work is the density and distribution of such confinement resonances and their possible relation to quantum chaos.

Because classical calculations of atom trajectories in comparable 1D systems have revealed signatures of chaos [19, 20], the statistical properties of the corresponding quantum mechanical resonances are of keen interest. According to a conjecture by Bohigas, Giannoni, and Schmit [29], distributions of the nearest-neighbor energy level spacings in the limit of strong chaos should obey the Wigner-Dyson (WD) distribution [30–33]. We investigate the structure of resonances that arise from atom-ion complex states from a fully quantum-mechanical calculation.

We propose a simple, one-dimensional model for the collision of a free ^{6}Li atom with a trapped ^{171}Yb$^+$ ion. We neglect trap micromotion, and assume harmonic confinement for the ion. We solve the Schrödinger equation for the reactance K-matrix, and compute both elastic and inelastic cross sections for the atom-ion collision, which can take the ion from one oscillator state to another.

This manuscript is arranged as follows. In Section II we detail the methods used in the work, including the