



Control of single spin-flips in a Rydberg atomic fractal

R. C. Verstraten[1]*, I. H. A. Knottnerus[2,3,4]*, Y. C. Tseng[2,3], A. Urech[2,3], T. S. do Espirito Santo[1,5], V. Zampronio[1,5,6], F. Schreck[2,3], R. J. C. Spreeuw[2,3], and C. Morais Smith[1#]

[1]*Institute for Theoretical Physics, Utrecht University, Princetonplein 5, 3584CC Utrecht, The Netherlands*
[2]*Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098XH Amsterdam, The Netherlands*
[3]*QuSoft, Science Park 123, 1098XG Amsterdam, The Netherlands*
[4]*Eindhoven University of Technology, P.O. Box 513, 5600MB Eindhoven, The Netherlands*
[5]*Departamento de Física Teórica e Experimental, Universidade Federal do Rio Grande do Norte, Campus Universitário, Lagoa Nova, Natal-RN 59078-970, Brazil*
[6]*Dipartimento di Fisica e Astronomia, Universitá di Firenze, I-50019, Sesto Fiorentino (FI), Italy*

Date: August 14 2025

* These authors contributed equally to this work
Corresponding author. Email: c.demoraissmith@uu.nl

Rydberg atoms trapped by optical tweezers have emerged as a versatile platform to emulate lattices with different geometries, in which long-range interacting spins lead to fascinating phenomena, ranging from spin liquids to topological states of matter. Here, we show that when the lattice has a fractal geometry with Hausdorff dimension 1.58, additional surprises appear. The system is described by a transverse-field Ising model with long-range van der Waals interactions in a Sierpiński gasket fractal. We investigate the problem theoretically using exact diagonalization, variational mean field, quantum Monte Carlo, and a graph-based numerical technique, SIM-GRAPH, which we developed. We find that in the quantum regime, the phase diagram exhibits phases in which the spins flip one-by-one. The theoretical results are in excellent agreement with experiments performed with single ^{88}Sr atoms trapped by optical tweezers arranged in a fractal geometry. The magnetization and von Neumann entanglement entropy reveal several regimes in which single spin-flips are delocalized over many sites of one sublattice, thus allowing for an unprecedented control of a cascade of phase transitions in a many-body system. These results expand the possibilities of Rydberg atoms for quantum information processing and may have profound implications in quantum technology.