



2022–
2024

HIGH-DIMENSIONAL QUANTUM SCHUR TRANSFORMS

ADAM BURCHARDT[1,2], JIANI FEI[3], DMITRY GRINKO[1,4,5], MARTIN LAROCCA[6], MARIS OZOLS[1,4,5], SYDNEY TIMMERMAN[3], AND VLADYSLAV VISNEVSKYI[1,7]

Abstract. The quantum Schur transform has become a foundational quantum algorithm, yet even after two decades since the seminal 2005 paper by Bacon, Chuang, and Harrow (BCH), some aspects of the transform remain insufficiently understood. Moreover, an alternative approach proposed by Krovi in 2018 was recently found to contain a crucial error. In this paper, we present a corrected version of Krovi's algorithm along with a detailed treatment of the high-dimensional version of the BCH Schur transform. This high-dimensional focus makes the two versions of the transform practical for regimes where the number of qudits n is smaller than the local dimension d, with Krovi's algorithm scaling as $\widetilde{O}(n^4)$ and BCH as $\widetilde{O}(\min(n^5, nd^4))$. Our work addresses a key gap in the literature, strengthening the algorithmic foundations of a wide range of results that rely on Schur–Weyl duality in quantum information theory and quantum computation.

arXiv:2509.22640v1 [quant-ph] 26 Sep 2025

Contents

1. Introduction 1
2. Preliminaries 3
3. Two approaches to the Schur transform 7
4. Revised Krovi's Schur transform 8
5. Revised Krovi's approach implements Gelfand–Tsetlin bases for S_n and U_d 11
6. Quantum circuits for step 1: preprocessing circuits 17
7. Quantum circuits for step 3: implementation of the isometry V 25
8. High-dimensional BCH Schur transform 29
Acknowledgements 32
References 32

1. Introduction

1.1. **Background.** The *Schur–Weyl duality* is a classic and fundamental result in representation theory. From an algebraic perspective, it is a theorem about two subalgebras of the full matrix algebra, one generated by the unitary group and another by the symmetric group. Specifically, Schur–Weyl duality says that the diagonal action of the unitary group U_d on the tensor space $\mathcal{H} := (\mathbb{C}^d)^{\otimes n}$ generates a matrix algebra that is the centralizer of the matrix algebra generated by the tensor representation of the symmetric group S_n, and vice versa. In this sense, Schur–Weyl duality is a consequence of the double centralizer theorem [Eti+11].

However, from a representation-theoretic perspective the Schur–Weyl duality presents a much deeper argument about relating finite-dimensional irreducible representations (irreps) of the unitary and the symmetric groups. In particular, it claims that the tensor space $\mathcal{H}$ decomposes into a direct sum of tensor products, each formed between a pair of irreducible modules: one of the unitary group and one of the symmetric group. Each such pair carries a shared irrep label, thus relating the irreps of the two groups.

Any basis in which the tensor space $\mathcal{H}$ decomposes into such a direct sum over tensor products of irreducible modules of the unitary and symmetric groups is called a *Schur basis*, and any unitary transformation from the standard computational basis to such basis is called a *Schur transform*.

One notable consequence of the decomposition in a Schur basis is that the symmetric group and the unitary group act on different modules, and thus on different registers in the tensor product pairs. This separation makes Schur transform a powerful tool for quantum algorithms and protocols that exploit either permutation or

(1) QuSoft, Amsterdam
(2) Centrum Wiskunde & Informatica, Amsterdam
(3) Stanford Institute for Theoretical Physics, Stanford University
(4) Institute for Logic, Language and Computation, University of Amsterdam
(5) Korteweg-de Vries Institute for Mathematics, University of Amsterdam
(6) Theoretical Division, Los Alamos National Laboratory
(7) Institute of Physics, University of Amsterdam
E-mail addresses: adam.burchardt.uam@gmail.com, jianif@stanford.edu, grinko.dimitry@gmail.com, laroccamartin@gmail.com, marozols@gmail.com, sydneyt@stanford.edu, vladislav.visnevskyy@gmail.com.