

2022–
2025

Optimizing fermionic Hamiltonians with classical interactions

Maarten E. Stroeks[1,2], Barbara M. Terhal[1,2], and Yaroslav Herasymenko[1,2,3,4]

[1]Delft Institute of Applied Mathematics, TU Delft, The Netherlands [2]QuTech, TU Delft, The Netherlands
[3]QuSoft & CWI, Amsterdam, The Netherlands [4]Perimeter Institute, Waterloo, Canada

arXiv:2510.02122v1 [quant-ph] 2 Oct 2025

Abstract

We consider the optimization problem (ground energy search) for fermionic Hamiltonians with classical interactions. This QMA-hard problem is motivated by the Coulomb electron-electron interaction being diagonal in the position basis, a fundamental fact that underpins electronic-structure Hamiltonians in quantum chemistry and condensed matter. We prove that fermionic Gaussian states achieve an approximation ratio of at least 1/3 for such Hamiltonians, independent of sparsity. This shows that classical interactions are sufficient to prevent the vanishing Gaussian approximation ratio observed in SYK-type models. We also give efficient semi-definite programming algorithms for Gaussian approximations to several families of traceless and positive-semidefinite classically interacting Hamiltonians, with the ability to enforce a fixed particle number. The technical core of our results is the concept of a Gaussian blend, a construction for Gaussian states via mixtures of covariance matrices.

1 Introduction

In this paper we study energy optimization, or ground energy search, for fermionic Hamiltonians. Mathematically, it means finding the largest[1] eigenvalue of a 2^n-dimensional Hermitian matrix, which is a low-degree polynomial in fermionic creation and annihilation operators $\{a_j^\dagger, a_j\}_{j\in[n]}$:

$$a_j^\dagger a_k + a_k a_j^\dagger = \delta_{jk}, \quad a_j a_k + a_k a_j = 0, \quad n_j \left|\boldsymbol{x}\right\rangle = x_j \left|\boldsymbol{x}\right\rangle, \tag{1}$$

where $n_j \equiv a_j^\dagger a_j$ and $\left|\boldsymbol{x}\right\rangle$ for $\boldsymbol{x} = (x_1, \ldots, x_n) \in \{0,1\}^n$ are the computational basis states. Energy optimization is one of the key computational problems in many-body physics and appears in a number of contexts in condensed matter physics and quantum chemistry. In general, it is a QMA-hard optimization task [1, 2]; in numerical practice, it is being accomplished with a number of approximation tools [3–7]. An interesting goal mathematically is to give rigorous performance guarantees for such approximate methods.

This text focuses on approximating the highest energy state with a Gaussian (i.e., free-fermionic) state [8–10]. Generally, Gaussian states are defined as the Gibbs states of Hamiltonians which are quadratic in $\{a_j^\dagger, a_j\}_{j\in[n]}$; they admit a classically efficient description in terms of a $2n$-sized *covariance matrix* (morally analogous to the stabilizer tableau for stabilizer states). In the domain of computational many-body physics, the standard method for finding Gaussian ground state approximations is generalized Hartree-Fock, which is heuristic [3, 4]. But in recent years, also *rigorous* guarantees for Gaussian ground state approximations (or lack thereof) have started to appear [11–18]. These use a common metric for any optimization method — *approximation ratio*, i.e., a guarantee on the ratio between the energy of the state obtained by a method, and the true ground energy[2]. It was discovered in [12, 13] that for *general* fermionic Hamiltonians, Gaussian states cannot yield ground energy even up to a constant approximation ratio. This dramatic effect – let us call it Gaussian (approximation) breakdown – was demonstrated for the Sachdev-Ye-Kitaev model. It was later extended to some other models which share the feature of having all-to-all, or at least non-sparse, fermion couplings [14, 16–18]. This breakdown can be viewed as a heuristic warning sign for optimization of general quantum chemistry Hamiltonians, as those are also

[1]We flip the sign convention of the Hamiltonian to align with that used in computer science.
[2]In the fermionic optimization literature, this ratio has been made well-defined by considering traceless Hamiltonians.