

2023–
2024







Is it Gaussian? Testing bosonic quantum states

Filippo Girardi[♣,1,*] Freek Witteveen[♣,2,†] Francesco Anna Mele,[3,‡] Lennart Bittel,[4,§] Salvatore F. E. Oliviero,[1,¶] David Gross,[5,**] and Michael Walter[6,7,8,††]

[1]*Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy*
[2]*QuSoft and Centrum Wiskunde & Informatica, Science Park 123, Amsterdam, the Netherlands*
[3]*NEST, Scuola Normale Superiore and Istituto Nanoscienze, Piazza dei Cavalieri 7, IT-56126 Pisa, Italy*
[4]*Dahlem Center for Complex Quantum Systems, Freie Universität Berlin, 14195 Berlin, Germany*
[5]*Institute for Theoretical Physics, University of Cologne, Zülpicher Straße 77, Cologne, Germany*
[6]*Ludwig-Maximilians-Universität München, Theresienstraße 37, Munich, Germany*
[7]*Ruhr University Bochum, Universitätsstraße 150, Bochum, Germany*
[8]*KdVI and QuSoft, University of Amsterdam, Science Park 123, Amsterdam, the Netherlands*

arXiv:2510.07305v1 [quant-ph] 8 Oct 2025

Gaussian states are widely regarded as one of the most relevant classes of continuous-variable (CV) quantum states, as they naturally arise in physical systems and play a key role in quantum technologies. This motivates a fundamental question: given copies of an unknown CV state, how can we efficiently test whether it is Gaussian? We address this problem from the perspective of representation theory and quantum learning theory, characterizing the sample complexity of Gaussianity testing as a function of the number of modes. For pure states, we prove that just a *constant* number of copies is sufficient to decide whether the state is *exactly* Gaussian. We then extend this to the tolerant setting, showing that a polynomial number of copies suffices to distinguish states that are close to Gaussian from those that are far. In contrast, we establish that testing Gaussianity of general mixed states necessarily requires exponentially many copies, thereby identifying a fundamental limitation in testing CV systems. Our approach relies on rotation-invariant symmetries of Gaussian states together with the recently introduced toolbox of CV trace-distance bounds.

I. INTRODUCTION

Property testing [1, 2] concerns the following broadly applicable scenario: you can sample from an unknown random variable, or have access to copies of an unknown quantum state, and you want to know whether the random variable or quantum state has some given property. Ideally, one would like to determine this with high confidence, but using as few samples or copies as possible with respect to the system size.

While property testing has been extensively studied in finite-dimensional quantum systems, its application to continuous-variable (CV) quantum systems—where states are defined over infinite-dimensional Hilbert spaces—remains largely unexplored. This gap is particularly striking given the recent surge of interest in quantum learning theory for CV systems [3–20]. A significant portion of this literature focuses on quantum state tomography, where the efficiency landscape is sharply divided: tomography of general non-Gaussian states is provably inefficient [3], while Gaussian states admit efficient procedures [3–6].

In this work, we address a fundamental question in this context: *Can we efficiently determine whether an unknown CV state is Gaussian?* We address a refined version of this question, where we determine the precise sample complexity of this task depending on the deviation from Gaussianity. This question is of intrinsic interest as a fundamental information-theoretic and statistical question on CV quantum systems, given the prominent role of Gaussian states in theory of CV quantum systems and their importance in quantum optics. It also has direct applications for CV-based quantum technologies. Gaussian states underpin a broad range of applications, including quantum metrology [21–23], quantum cryptography [24–27], universal quantum computation [28], and Gaussian boson sampling [29–31]. A robust test for Gaussianity is therefore an valuable tool for a wide range of applications. For example, if one can reliably identify a state or process as Gaussian, one can then apply specialized tomography algorithms to learn the full state [3–6].

Previous works on property testing have addressed the problem of testing stabilizer states [32–35] and fermionic Gaussian states [36, 37]. A common theme with these works is that good property tests can be derived from symmetry considerations. We will see that this approach can be made to work in the continuous-variable setting of testing bosonic Gaussian states as well.

A. Summary of results

We now present an overview of our results. In order to test whether a given state is Gaussian, we must use some condition which singles out Gaussian states compared to other states. We now explain two conditions characterizing Gaussianity, and explain in more detail

♣ These authors contributed equally to this work.