

Anti-concentration is (almost) all you need

Markus Heinrich,[1,*] Jonas Haferkamp,[2,3] Ingo Roth,[4] and Jonas Helsen[5]

[1] *Institute for Theoretical Physics, University of Cologne, Germany*
[2] *School of Engineering and Applied Science, Harvard University, USA*
[3] *Department of Mathematics, Saarland University, Germany*
[4] *Quantum Research Center, Technology Innovation Institute, Abu Dhabi, United Arab Emirates*
[5] *QuSoft and CWI, Amsterdam, Netherlands*

Until very recently, it was generally believed that the (approximate) 2-design property is strictly stronger than anti-concentration of random quantum circuits, mainly because it was shown that the latter anti-concentrate in logarithmic depth, while the former generally need linear depth circuits. This belief was disproven by recent results which show that so-called *relative-error* approximate unitary designs can in fact be generated in logarithmic depth, implying anti-concentration. Their result does however not apply to ordinary local random circuits, a gap which we close in this letter, at least for 2-designs. More precisely, we show that anti-concentration of local random quantum circuits already implies that they form relative-error approximate state 2-designs, making them equivalent properties for these ensembles. Our result holds more generally for any random circuit which is invariant under local (single-qubit) unitaries, independent of the architecture.

arXiv:2510.23719v2 [quant-ph] 8 Dec 2025

Random quantum circuits are ubiquitous in quantum information theory, touching a wide range of topics from randomized benchmarking [1–5] to black hole physics [6, 7]. Consequently, significant effort has been devoted to studying the mixing properties of random quantum circuits.

A particular focus of the community concerns the convergence of random quantum circuits to approximate (unitary or state) k-designs. These are probability distributions on the unitary group or state space that appear uniformly random given access to at most k copies. Early works rigorously proved the convergence of random quantum circuits to approximate 2-designs in depth $O(n)$ on n qubits [8, 9]. The convergence was subsequently tied to spectral properties of the k-copy twirling channel [10, 11], although tight bounds were still limited to $k = 2$. Since then, a long line of works extended this result to higher k [12–15], ultimately providing near tight bounds on the k-dependence [16] in this convergence.

At the same time, the linear dependence on the system size was widely considered optimal. This however changed very recently, when it was shown that random quantum circuits already look Haar-random to forward-in-time experiments at logarithmic depth [17, 18].

In hindsight, the rapid onset of *anti-concentration* [19, 20] could be considered as a first hint for the logarithmic convergence of second moments. Anti-concentration of a circuit ensemble ν states, loosely speaking, that the distribution of measurement outcomes obtained from such circuits is typically 'well spread'. This property is a key ingredient in complexity-theoretic arguments for sampling-based quantum advantage (see Ref. [21] for a recent review). Anti-concentration is implied by a small value of the (average) *collision probability*

$$Z_\nu = \mathbb{E}_{U\sim\nu}|\langle 0|U|0\rangle|^4 . \tag{1}$$

Using a mapping to a statistical mechanics model [22–24], it was shown in Refs. [19, 20] that random quantum circuits with iid gates have a near minimal average collision probability already at log-depth.

The logarithmic-depth designs constructed in Refs. [17, 18] can be seen as 'coarse-grained' brickwork circuits in the sense that certain entangling blocks are deleted. This raises the question whether this deletion is actually advantageous, or whether it is simply a remnant of the proof technique. Intuitively, the insertion of random gates into the circuit should not slow down the mixing time, but such effects are known to occur in Markov chains. A similar behavior was recently observed in quantum circuits involving Haar-random unitaries that act on an extensive number of qubits [25]. As such, the latter does however not extend to local random quantum circuits.

In this letter, we provide the first evidence that standard random quantum circuit indeed converge as fast as the coarse-grained circuits in Refs. [17, 18] by proving the convergence of standard brickwork random quantum circuits to so-called relative-error approximate *state* 2-designs in logarithmic depth. In other words, we show that in the case of the state 2-design property the extra structure in Refs. [17, 18] is not necessary and that the result of Ref. [25] does not qualitatively apply to local random quantum circuits. Our proof is remarkably short and reduces the relative-error state 2-design property to the convergence of the collision probability (i.e. *anti-concentration*). Interestingly, the inverse implication is always true [26], hence *anti-concentration is all you need* for the here considered state ensembles. Finally, we also briefly discuss the case of unitary designs and why anti-concentration may generally not be enough there. These intricacies are illustrated in parallel work [27] which also gives evidence that anti-concentration may nevertheless be enough for the unitary 2-design property in 1D brickwork circuits.

Complementary evidence for the fast convergence of