



Engineering Zeeman-manifold quintets using state-dependent light shifts in neutral atoms

Benedikt Heizenreder [1, *] Bas Gerritsen [2, 3, *] Katya Fouka [2, 3, *] Robert J. C. Spreeuw [1, 3] Florian Schreck [1, 3] Arghavan Safavi Naini [2, 3] and Alexander Urech [1, 3, †]

[1] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

[2] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

[3] *QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*

(Dated: December 17, 2025)

We present a general method for engineering qudits through individually addressable transitions between Zeeman sublevels, achieved by combining a large linear Zeeman shift with a state-dependent light shift. This approach lifts the degeneracy between adjacent states while simultaneously tuning their energy splittings into the radio-frequency (RF) domain, enabling coherent manipulation within the Zeeman manifold using experimentally accessible drive frequencies. As a concrete realization, we investigate the implementation of an $SU(5)$ *quintet* encoded in the Zeeman sublevels of the long-lived $^3P_2$ state of neutral ^{88}Sr atoms confined in far-detuned, σ^--polarized optical tweezers. Using realistic experimental parameters, we numerically demonstrate full control of the *quintet* manifold, including initialization into a specific $SU(5)$ basis state via a multi-photon transfer, coherent state- and site-selective single-qudit rotations driven by RF fields, and fast state-selective optical readout. Our simulations predict state-preparation fidelities of $\mathcal{F} \simeq 0.99$ within ~ 1 µs, single-qudit gate fidelities of $\mathcal{F} \simeq 0.99$ with π-pulse durations of ~ 2.5 µs, and fast destructive imaging with durations below 10 µs. These results establish a broadly applicable framework for high-fidelity control of Zeeman sublevel-encoded qudits and highlight the $^3P_2$ manifold in strontium as a promising platform for scalable qudit-based quantum technologies.

arXiv:2512.14611v1 [physics.atom-ph] 16 Dec 2025

I. INTRODUCTION

Neutral alkaline-earth-metal atoms trapped in arrays of optical tweezers or optical lattices form a powerful platform for atomic clocks and sensors [1–3], quantum computing [4–6], and quantum simulation of many-body physics [7–9]. Their success is largely driven by the coherent control of optical intercombination transitions, exploiting extremely long-lived excited states that enable coherence times on the order of minutes [10]. At the same time, these advantages require highly coherent and long-term stable laser systems [11, 12]. Moreover, the need for multiple fast rotations and sophisticated decoupling sequences in the optical domain remains a significant experimental challenge [13, 14].

Parallel to these developments, there is a growing interest in exploiting higher-dimensional Hilbert spaces, i.e. $SU(N > 2)$ physics [15–18]. Coherent control over multiple internal atomic levels enables the realization of so-called *qudits*. Qudits are not only of interest for quantum simulation but also could offer significant advantages for quantum computing, including improvements in quantum error correction and scalable fault-tolerant architectures [19, 20]. Furthermore, they reduce physical resource requirements by enabling the encoding of logical qubits within a single atom [21–24].

One can use different approaches to realize such qudits in neutral atoms. One strategy encodes information in nuclear spins of the electronic ground state, utilizing their exceptional isolation from the environment and long coherence times [25–28]. Alternatively, metastable electronic sub-levels [21, 29–31] or even a combination of nuclear spins and electronic levels provide attractive opportunities [32]. Particularly appealing are qudits where computational coupling is performed in the radio-frequency (RF) domain [21, 29, 33]. The long RF wavelength enables phase-coherent control, precise and fast rotations, and the implementation of demanding dynamical decoupling sequences [34–36]. These operations are relatively easy to implement compared to the optical domain. However, while high-fidelity control in the RF domain is well established, preparation and readout of the qudit must still be performed optically, and scaling such methods to systems with more than two levels remains a key challenge [27, 29, 30].

In this work, we study the realization of an $SU(5)$ *quintet* encoded in neighboring Zeeman sublevels. To lift the degeneracy between these levels, we combine a large linear Zeeman shift, induced by an external magnetic field, with a state-dependent light shift generated by the σ-polarized optical tweezer beams (see Fig. 1). This approach engineers non-degenerate energy splittings between adjacent states, allowing individual RF drive frequencies within a range easily accessible to standard RF technology (~ 100 MHz). We specifically explore a potential experimental realization (Sec. II A) that benefits from the long lifetime (tens of seconds) of the $^3P_2$ state in

* These authors contributed equally to this work.
† Quintet3P2@strontiumBEC.com