


Direct observation of the optical Magnus effect with a trapped ion

Philip Leindecker,[1,2,*] Louis P.H. Gallagher,[3,†] Edgar Brucke,[1,2] Dominique Zehnder,[1,2] Luka Milanovic,[1,2] Matteo Marinelli,[1,2,4] Rene Gerritsma,[3,5] Robert J.C. Spreeuw,[3,5] Jonathan Home,[2,6] and Cornelius Hempel[1,2,6]

[1] *ETH Zürich – PSI Quantum Computing Hub, Paul Scherrer Institut (PSI), 5232 Villigen, Switzerland*
[2] *Institute for Quantum Electronics, ETH Zürich, Otto-Stern-Weg 1, 8093 Zürich, Switzerland*
[3] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, 1098 XH Amsterdam, Netherlands*
[4] *Department of Physics, University of Trieste, 34127 Trieste, Italy*
[5] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[6] *Quantum Center, ETH Zurich, CH-8093, Switzerland*

(Dated: February 2, 2026)

We directly observe and spatially map an optical analog of the Magnus effect, where intrinsic spin-orbit-like coupling of light generates a spin-dependent transverse displacement of the atom–light interaction profile for a $^{40}Ca^{+}$ ion. Probed on a quadrupole transition using a tightly focused beam, we observe displacements of the maximum in the profile of the effective interaction by several 100 nm originating from intrinsic longitudinal electric field components beyond the paraxial approximation. The tight focus of the beam induces additional transverse polarization gradients, which we characterize through a phase-sensitive measurement and spatial maps for different beam configurations. The results establish the physical basis of polarization-gradient interactions relevant to optical tweezer-based quantum control.

arXiv:2601.22981v1 [physics.atom-ph] 30 Jan 2026

The Magnus effect [1] describes how a rotating object moving through a medium experiences a force that is orthogonal to both its velocity and its axis of rotation. In sports, this effect is widely exploited to send a spinning ball along a curved trajectory, controlled by the direction of rotation. An optical analog of the Magnus effect has been predicted for atoms trapped in optical tweezers [2, 3], where intrinsic spin-orbit-like coupling of light (Appendix A) gives rise to a spin dependent transverse structure of the atom-light interaction, which is not present in the plane wave approximation.

A tightly focused Gaussian beam without intrinsic orbital angular momentum (OAM) acquires local OAM and longitudinal electric field components in the nonparaxial description. The interplay of the induced OAM and the beam's spin angular momentum produces spin-orbit-like coupling and an inverted wavefront curvature across the focal plane, generating strong transverse polarization gradients and shifting the maxima of its circular field components off axis [4–8]. When an atom is placed within the beam, the shift of these field components leads to a transverse displacement of the effective atom-light interaction profile. The predicted shift is wavelength (λ) dependent and calculated to lie between $\lambda/2\pi$ and λ/π for different components of the quadrupole transition [9] and $\lambda/2\pi$ for dipole transitions [2, 3]. At the beam center, this shift causes the atom to experience an intensity gradient of the light, thereby enabling transverse motional excitations forbidden for a paraxial Gaussian beam, analogous to the interaction of an atom with structured light containing intrinsic OAM [10–15]. A related polarization dependent spatial dependence of the origin of displacement in light *emission* was previously observed with a trapped ion [16].

Spin-dependent transverse forces can introduce unwanted coupling to the motion of ions positioned at the center of the beam [9, 17–19], but the same mechanism enables new optical quantum gate schemes based on engineered polarization gradients [20–22]. As optical tweezers become increasingly central to atomic physics – enabling parallel trapping and control of large neutral-atom arrays [23–28] and offering new routes to high-fidelity trapped-ion gates [20, 29–32] – understanding such intrinsic transverse interactions becomes important.

In this Letter, we report the first direct observation of the optical Magnus effect in the interaction of a single trapped ion with a tightly focused Gaussian beam. Using sub-100-nm spatial resolution in the positioning of laser beams enabled by two crossed acousto-optic deflectors (AODs) [33], we map the spatial profile of coupling strength of all components of the 729 nm quadrupole transition from $4S_{1/2} \rightarrow 3D_{5/2}$ in $^{40}Ca^{+}$. We resolve the spin-dependent transverse displacement of the interaction profile arising solely from the Gaussian beam's intrinsic longitudinal components [2, 3]. We also characterize additional transverse polarization gradients arising from the tight focus of the beam leading to position-dependent selection rules for the quadrupole transition.

The light-matter interaction can be formally described as follows: the electric field of the tightly focused tweezer beam can be approximated to first order beyond the paraxial limit as [34, 35]

$$\mathbf{E}(\mathbf{r},t) \approx \mathrm{Re}\left[\left(\hat{\boldsymbol{\varepsilon}} - \frac{i\,(\varepsilon_x x + \varepsilon_y y)}{z_0}\hat{\mathbf{z}}\right) f(\mathbf{r}) e^{i(kz-\omega t)}\right], \quad (1)$$

where $\hat{\boldsymbol{\varepsilon}} = \varepsilon_x\hat{\mathbf{x}} + \varepsilon_y\hat{\mathbf{y}}$ is the transverse polarization unit vector, $z_0 = \pi w_0^2/\lambda = k w_0^2/2$ is the Rayleigh range and $f(\mathbf{r})$ is the Gaussian beam mode function, defined explicitly alongside the other parameters in Appendix B. The Hamiltonian of the quadrupole atom-light interaction is