

Plethysm is in #BQP

Matthias Christandl[1], Aram W. Harrow[2], Greta Panova[3], Pietro M. Posta[1], and Michael Walter[4,5,6]

[1]Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen, Denmark
[2]Center for Theoretical Physics – a Leinweber Institute, MIT, Cambridge, MA, 02139, USA
[3]Department of Mathematics, University of Southern California, Los Angeles, CA, 90089, USA
[4]Ludwig-Maximilians-Universität München, Theresienstr. 37, 80333 München, Germany
[5]Munich Center for Quantum Science and Technology (MCQST), Schellingstr. 4, 80799 München, Germany
[6]Korteweg-de Vries Institute for Mathematics & QuSoft, University of Amsterdam, The Netherlands

arXiv:2602.08441v1 [quant-ph] 9 Feb 2026

Abstract

Some representation-theoretic multiplicities, such as the Kostka and the Littlewood-Richardson coefficients, admit a combinatorial interpretation that places their computation in the complexity class #P. Whether this holds more generally is considered an important open problem in mathematics and computer science, with relevance for geometric complexity theory and quantum information. Recent work has investigated the quantum complexity of particular multiplicities, such as the Kronecker coefficients and certain special cases of the plethysm coefficients.

Here, we show that a broad class of representation-theoretic multiplicities is in #BQP. In particular, our result implies that the plethysm coefficients are in #BQP, which was only known in special cases. It also implies all known results on the quantum complexity of previously studied coefficients as special cases, unifying, simplifying, and extending prior work. We obtain our result by multiple applications of the Schur transform. Recent work has improved its dependence on the local dimension, which is crucial for our work. We further describe a general approach for showing that representation-theoretic multiplicities are in #BQP that captures our approach as well as the approaches of prior work. We complement the above by showing that the same multiplicities are also naturally in GapP and obtain polynomial-time classical algorithms when certain parameters are fixed.

1 Introduction

Computing multiplicities is a central problem in representation theory and algebraic combinatorics, and is directly relevant to quantum information theory and geometric complexity theory. In most cases, no closed-form formulas are known. A natural and fruitful question is to ask whether these multiplicities count combinatorial objects. If so, then one obtains *positive combinatorial interpretation*. Stanley stated the problem of finding such interpretations for important representation-theoretic multiplicities in his influential paper [54]. Specific multiplicities of interest include: the *Littlewood-Richardson (LR) coefficients*, discussed in 1934 [39], giving the multiplicity of an irreducible representation of the general linear group $\mathrm{GL}(n)$ in the tensor product of two others; the *Kronecker coefficients* of the symmetric group, defined in 1938 [42], giving the multiplicity of an irreducible representation of the symmetric group S_n in the tensor product of two others; and the *plethysm coefficients*, first mentioned by Littlewood in 1936 [37] and defined formally in Ref. [38], which gives the multiplicity of an irreducible $\mathrm{GL}(n)$-representation in the composition of two irreducible GL-representations (see Section 1.3). By Schur-Weyl duality, the