

Optimal learning of covariant quantum states and channels

Satoshi Yoshida[*], Kazuki Okigami[†], Pietro M. Posta[‡], Dmitry Grinko[§]

October 1, 2026

arXiv:2609.39280v1 [quant-ph] 30 Sep 2026

Abstract

We give collective tomography protocols for quantum states and channels with known symmetries, using random purification and dilation to reduce learning to pure-state estimation. For states commuting with a compact-group representation with multiplicities m_λ, the optimal copy complexity is $\Theta((\sum_\lambda m_\lambda^2 + \log \eta^{-1})/\varepsilon^2)$ for sufficiently small trace-distance error ε and failure probability η for the nontrivial case $\sum_\lambda m_\lambda^2 > 1$. For G-covariant channels with finite-dimensional unitary representations of a compact group G, parallel $\widetilde{O}(D_G/\varepsilon^2)$ queries achieve a diamond-distance error ε at fixed success probability $2/3$, where D_G counts the real parameters of covariant Choi operators before imposing trace preservation. For permutation-covariant channels on k qudits of fixed local dimension d, at sufficiently small fixed error and fixed success probability $2/3$, we obtain the optimal query scalings $\Theta_d(k^{d^4-1})$ for diamond distance and $\Theta_d(k^{d^4-d^2})$ for Choi trace distance. Furthermore, we resolve an open problem in quantum tomography of constructing efficient quantum circuits that approximate the Hayashi measurement for optimal pure-state estimation. We encode states in the symmetric subspace into bosonic occupation modes, realize the measurement by heterodyne detection and normalization, and approximate this procedure on qubits using the quantum Hermite transform. Combined with our symmetry-compatible purification and dilation circuits, this yields query-optimal and gate-efficient learners of permutation-covariant states and channels with gate complexity $O_d(\mathrm{poly}(k, \varepsilon^{-1}, \log \eta^{-1}))$.

[*]Department of Physics, Graduate School of Science, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan. satoshiyoshida.phys@gmail.com

[†]OptQC Corp., 3-28-13 Nishi-Ikebukuro, Toshima-ku, Tokyo 171-0021, Japan. kazuki.okigami@optqc.com

[‡]Department of Mathematical Sciences, University of Copenhagen, Denmark. pmp@math.ku.dk

[§]QuSoft & Institute for Logic, Language and Computation & Korteweg-de Vries Institute for Mathematics, University of Amsterdam, The Netherlands. d.grinko@uva.nl