

2017–2021

arXiv:1609.06994v2 [quant-ph] 12 Oct 2018

Deconstruction and conditional erasure of quantum correlations

Mario Berta
Department of Computing, Imperial College London, London SW7 2AZ, UK

Fernando G. S. L. Brandão
Institute for Quantum Information and Matter, California Institute of Technology, Pasadena, California 91125, USA

Christian Majenz
Institute for Language, Logic and Computation, University of Amsterdam, and QuSoft, 1098XG Amsterdam, Netherlands and
Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, DK-2100 Copenhagen Ø

Mark M. Wilde
Hearne Institute for Theoretical Physics, Department of Physics and Astronomy, Center for Computation and Technology, Louisiana State University, Baton Rouge, Louisiana 70803, USA

(Dated: October 15, 2018)

We define the deconstruction cost of a tripartite quantum state on systems ABE as the minimum rate of noise needed to apply to the AE systems, such that there is negligible disturbance to the marginal state on the BE systems, while the system A of the resulting state is locally recoverable from the E system alone. We refer to such actions as deconstruction operations and protocols implementing them as state deconstruction protocols. State deconstruction generalizes Landauer erasure of a single-party quantum state as well the erasure of correlations of a two-party quantum state. We find that the deconstruction cost of a tripartite quantum state on systems ABE is equal to its conditional quantum mutual information (CQMI) $I(A;B|E)$, thus giving the CQMI an operational interpretation in terms of a state deconstruction protocol. We also define a related task called conditional erasure, in which the goal is to apply noise to systems AE in order to decouple system A from systems BE, while causing negligible disturbance to the marginal state of systems BE. We find that the optimal rate of noise for conditional erasure is also equal to the CQMI $I(A;B|E)$. State deconstruction and conditional erasure lead to operational interpretations of the quantum discord and squashed entanglement, which are quantum correlation measures based on the CQMI. We find that the quantum discord is equal to the cost of simulating einselection, the process by which a quantum system interacts with an environment, resulting in selective loss of information in the system. The squashed entanglement is equal to half the minimum rate of noise needed for deconstruction/conditional erasure if Alice has available the best possible system E to help in the deconstruction/conditional erasure task.

I. INTRODUCTION

The Landauer erasure principle represents a deep link between information theory and thermodynamics [1]. An informal summary of the principle is that the work cost of erasing the contents of a computer memory is proportional to the amount of information stored there. This insight has now sparked a whole literature, a consequence of which has been a deepening of the connection between information theory and thermodynamics (see, e.g., [2] for a review).

One generalization of Landauer's insight goes beyond the single-system setup mentioned above. In [3], Groisman *et al.* considered a setting in which two parties share a quantum state ρ_{AB}. Their goal was to determine the work cost of erasing the correlations present in the state, by acting locally on one system, such that the resulting state has a tensor-product form $\sigma_A \otimes \omega_B$, where σ_A and ω_B are quantum states. Groisman *et al.* solved the problem in the framework of quantum Shannon theory [4], whereby they allowed the two parties to have many copies of the state ρ_{AB} and quantified the minimum rate of noise that needs to be applied to the A systems such that the resulting state is tensor-product between the A systems and the B systems. They found that the optimal rate of noise is equal to the quantum mutual information of the state ρ_{AB}, defined as

$$I(A;B)_\rho \equiv H(A)_\rho + H(B)_\rho - H(AB)_\rho, \tag{1}$$

where the quantum entropy of a state σ_G on system G is defined as $H(G)_\sigma \equiv -\operatorname{Tr}\{\sigma_G \log_2 \sigma_G\}$. An important consequence of their theorem is that we can assign a physical meaning to, or operational interpretation of, the quantum mutual information as the minimum rate of noise needed to completely erase the correlations present in a two-party quantum state. Thus, we can say that quantum mutual information is equal to the work cost of correlation destruction.

On the other hand, quantum mutual information has also been interpreted in a communication-theoretic task (now called *coherent state merging* [5]) as the optimal rate of entanglement creation when transferring the system A of ρ_{AB} to a party possessing system B [6], while using quantum communication at a fixed rate. These dual

