

Schur-Weyl duality for the Clifford group with applications: Property testing, a robust Hudson theorem, and de Finetti representations

David Gross* Sepehr Nezami† Michael Walter‡,†

arXiv:1712.08628v3 [quant-ph] 16 Jan 2021

Abstract

Schur-Weyl duality is a ubiquitous tool in quantum information. At its heart is the statement that the space of operators that commute with the t-fold tensor powers $U^{\otimes t}$ of all unitaries $U \in U(d)$ is spanned by the permutations of the t tensor factors. In this work, we describe a similar duality theory for tensor powers of *Clifford unitaries*. The Clifford group is a central object in many subfields of quantum information, most prominently in the theory of fault-tolerance. The duality theory has a simple and clean description in terms of finite geometries. We demonstrate its effectiveness in several applications:

- We resolve an open problem in *quantum property testing* by showing that "stabilizerness" is efficiently testable: There is a protocol that, given access to six copies of an unknown state, can determine whether it is a stabilizer state, or whether it is far away from the set of stabilizer states. We give a related membership test for the Clifford group.
- We find that tensor powers of stabilizer states have an increased symmetry group. Conversely, we provide corresponding *de Finetti theorems*, showing that the reductions of arbitrary states with this symmetry are well-approximated by mixtures of stabilizer tensor powers (in some cases, exponentially well).
- We show that the distance of a pure state to the set of stabilizers can be lower-bounded in terms of the sum-negativity of its Wigner function. This gives a new quantitative meaning to the sum-negativity (and the related *mana*) – a measure relevant to fault-tolerant quantum computation. The result constitutes a *robust* generalization of the *discrete Hudson theorem*.
- We show that complex projective designs of arbitrary order can be obtained from a finite number (independent of the number of qudits) of Clifford orbits. To prove this result, we give explicit formulas for arbitrary moments of random stabilizer states.

*Institute for Theoretical Physics, University of Cologne, and Centre for Engineered Quantum Systems, School of Physics, The University of Sydney, david.gross@thp.uni-koeln.de. †Stanford Institute for Theoretical Physics, Stanford University, and Kavli Institute for Theoretical Physics, nezami@stanford.edu. ‡Korteweg-de Vries Institute for Mathematics, Institute of Theoretical Physics, and Institute for Logic, Language and Computation, QuSoft, University of Amsterdam, m.walter@uva.nl.