

SciPost Physics Submission

Adiabatic state distribution using anti-ferromagnetic spin systems

K. Groenland

University of Amsterdam, QuSoft and Centrum Wiskunde en Informatica (CWI), Amsterdam, The Netherlands
k.l.groenland@cwi.nl

December 4, 2018

arXiv:1809.08158v2 [quant-ph] 5 Dec 2018

Abstract

Transporting quantum information is an important prerequisite for quantum computers. We study how this can be done in Heisenberg-coupled spin networks using adiabatic control over the coupling strengths. We find that qudits can be transferred and entangled pairs can be created between distant sites of bipartite graphs with a certain balance between the maximum spin of both parts, extending previous results that were limited to linear chains. The transfer fidelity in a small star-shaped network is numerically analysed, and possible experimental implementations are discussed.

Contents

1 Introduction **3**
1.1 Relation to previous work 3
1.2 Document structure 5

2 Ground states of symmetry-protected subspaces **5**
2.1 Preliminaries 5
2.2 Preservation of the ground state of $V_{s,m}$ 7

3 Applications **9**
3.1 Sharing a quantum state between multiple parties, such that any party can access the state 10
3.2 Distributing maximally entangled singlet states 11

4 Errors and scaling **12**
4.1 Numerics on star graphs 13
4.2 A numerical example where assumptions are violated 15

5 Experimental implementations **15**

6 Discussion and outlook **16**

7 Conclusion **17**

