

A Quantum Multiparty Packing Lemma and the Relay Channel

Dawei Ding, Hrant Gharibyan, Patrick Hayden, and Michael Walter

arXiv:1809.10149v2 [quant-ph] 26 May 2020

***Abstract*—Optimally encoding classical information in a quantum system is one of the oldest and most fundamental challenges of quantum information theory. Holevo's bound places a hard upper limit on such encodings, while the Holevo-Schumacher-Westmoreland (HSW) theorem addresses the question of how many classical messages can be "packed" into a given quantum system. In this article, we use Sen's recent quantum joint typicality results to prove a one-shot multiparty quantum packing lemma generalizing the HSW theorem. The lemma is designed to be easily applicable in many network communication scenarios. As an illustration, we use it to straightforwardly obtain quantum generalizations of well-known classical coding schemes for the relay channel: multihop, coherent multihop, decode-forward, and partial decode-forward. We provide both finite blocklength and asymptotic results, the latter matching existing classical formulas. Given the key role of the classical packing lemma in network information theory, our packing lemma should help open the field to direct quantum generalization.**

***Index Terms*—quantum channels, network coding, packing lemma, relay channel, simultaneous decoder**

I. Introduction

The *packing lemma* [1], [2], [3] is one of the central tools used in the construction and analysis of information transmission protocols [4]. It quantifies the asymptotic rate at which messages can be "packed" reversibly into a medium, in the sense that the probability of a decoding error vanishes in the limit of large blocklength. For concreteness, consider the following general version of the packing lemma.[1]

Lemma 1 (Classical Packing Lemma)**.** *Let (U, X, Y) be a triple of random variables with joint distribution p_{UXY}. For each n, let $(\tilde{U}^n, \tilde{Y}^n)$ be a pair of arbitrarily distributed random sequences and $\{\tilde{X}^n(m)\}$ a family of at most 2^{nR} random sequences such that each $\tilde{X}^n(m)$ is conditionally independent of $\tilde{Y}^n$ given $\tilde{U}^n$ (but arbitrarily dependent on the other $\tilde{X}^n(m')$ sequences). Further assume that each $\tilde{X}^n(m)$ is distributed as $\otimes_{i=1}^n p_{X|U=\tilde{U}_i}$ given $\tilde{U}^n$. Then, there exists $\delta(\varepsilon)$ that tends to zero as $\varepsilon \to 0$ such that*

$$\lim_{n\to\infty} \Pr((\tilde{U}^n, \tilde{X}^n(m), \tilde{Y}^n) \in \mathcal{T}_\varepsilon^{(n)} \text{ for some } m) = 0$$

if $R < I(X;Y|U) - \delta(\varepsilon)$, where $\mathcal{T}_\varepsilon^{(n)}$ is the set of ε-typical strings of length n with respect to p_{UXY}.

The packing lemma provides a unified approach to many, if not most, of the achievability results in Shannon theory. Despite its broad utility, it is a simple consequence of the union bound and the standard joint typicality lemma with the three variables U, X, Y. The usual channel coding theorem directly follows from taking $U = \emptyset$ and when $\tilde{Y}^n \sim p_Y^{\otimes n}$.

For the case when $U = \emptyset$ and when $\tilde{Y}^n \sim p_Y^{\otimes n}$, the quantum generalization of the packing lemma is known: the Holevo-Schumacher-Westmoreland (HSW) theorem [5], [6]. This can be proven using a conditional typicality lemma for a classical-quantum state with one classical and one quantum system. However, until recently no such typicality lemma was known for the case of multiple encoding systems, and so a quantum version of Lemma 1 was lacking. Furthermore, while in classical Shannon theory Lemma 1 can be used repeatedly in scenarios where the message is encoded into multiple random variables, this approach fails in the quantum case due to measurement disturbance, specifically the influence of one decoding on subsequent decodings. Hence, while it is sufficient to solve the full multiparty packing problem in the classical case with just two encoding systems and repeated measurements, a general multiparty packing lemma with $k \in \mathbb{N}$ encoding systems is required in the quantum case. The bottleneck is again the lack of a general quantum joint typicality lemma with multiple systems. However, we can obtain partial results in the quantum case for some network scenarios, as we will describe below.

In this paper we use the quantum joint typicality lemma[2] established recently by Sen [7] to prove a quantum one-shot multiparty packing lemma for k classical encoding systems. We then demonstrate the wide applicability of the lemma by using it to generalize classical network information theory protocols to the quantum case. The lemma allows us to construct and prove the correctness of these simple generalizations and, we believe, should help to open the field of classical network information theory to direct quantum generalization.[3] One feature of the lemma is that it leads naturally to demonstrations of the achievability of rate regions without having to resort to time-sharing, a desirable property known as *simultaneous decoding*. Simultaneous decoding is



Dawei Ding, Hrant Gharibyan, and Patrick Hayden are with the Stanford Institute for Theoretical Physics, Stanford University, Stanford, CA 94305, USA. Michael Walter is with QuSoft, the Korteweg-de Vries Institute for Mathematics, the Institute for Theoretical Physics, and the Institute for Logic, Language and Computation, University of Amsterdam, 1098 XG Amsterdam, The Netherlands. (c) 2018 IEEE. Personal use of this material is permitted. However, permission to use this material for any other purposes must be obtained from the IEEE by sending a request to pubs-permissions@ieee.org.

[1] See, e.g., [4]. Our formulation is slightly paraphrased and uses a notation that is more suitable for the following.

[2] Sen modestly calls his result a lemma, but the highly ingenious proof more than justifies calling it a theorem.

[3] Note that a simultaneous smoothing result for the max-relative entropy is still missing, which would be necessary e.g. for a "multiparty covering lemma." To prove such a result is a major open problem in the field.