

Quasirandom quantum channels

Tom Bannink[1], Jop Briët[2], Farrokh Labib[1], and Hans Maassen[3]

[1]CWI, QuSoft, Science Park 123, 1098 XG Amsterdam, Netherlands. Supported by the Gravitation-grant NETWORKS-024.002.003 from the Dutch Research Council (NWO).

[2]CWI, QuSoft, Science Park 123, 1098 XG Amsterdam, Netherlands. Supported by the Gravitation-grant NETWORKS-024.002.003 from the Dutch Research Council (NWO). Additionally supported by an NWO VENI grant

[3]QuSoft, Korteweg-de Vries Institute for Mathematics, Radboud University.

arXiv:1908.06310v3 [quant-ph] 8 Jul 2020

Mixing (or quasirandom) properties of the natural transition matrix associated to a graph can be quantified by its distance to the complete graph. Different mixing properties correspond to different norms to measure this distance. For dense graphs, two such properties known as spectral expansion and uniformity were shown to be equivalent in seminal 1989 work of Chung, Graham and Wilson. Recently, Conlon and Zhao extended this equivalence to the case of sparse vertex transitive graphs using the famous Grothendieck inequality.

Here we generalize these results to the non-commutative, or 'quantum', case, where a transition matrix becomes a quantum channel. In particular, we show that for irreducibly covariant quantum channels, expansion is equivalent to a natural analog of uniformity for graphs, generalizing the result of Conlon and Zhao. Moreover, we show that in these results, the non-commutative and commutative (resp.) Grothendieck inequalities yield the best-possible constants.

1 Introduction

In a seminal work [8], Chung, Graham and Wilson — building on work of Thomason [33, 34] — proved that several seemingly distinct notions of quasirandomness for graphs are equivalent. In particular, they identified seven properties found in random graphs with high probability, that always coexist simultaneously in any large dense graph. Two of these properties are *spectral expansion* and *uniformity* (defined below). A question of Chung and Graham [7] on the equivalence of these two properties in *sparse* graphs resulted in a line of research culminating in recent work of Conlon and Zhao [9], which introduced a surprising new item to the armory of combinatorics: the famous Grothendieck inequality [13]. In this paper, we draw a parallel line in the context of quantum information theory, where quantum channels take the place of graphs. In addition, we give a streamlined proof of the main result of [9] and show that the use of Grothendieck's inequality yields an optimal constant. Similarly, we show that the non-commutative Grothendieck inequality gives an optimal constant in the quantum setting.

Spectral expansion and uniformity. Spectral expansion is a linear-algebraic property given in terms of the transition matrix of a graph. This transition matrix is the normalized adjacency matrix, which for a d-regular graph $G = (V, E)$ is given by $A_{uv} = e(\{u\}, \{v\})/d$, where $e(S, T)$ denotes the number of edges connecting subsets $S, T \subseteq V$. We say that the graph G is an (n, d, λ) graph if $|V| = n$, it is d-regular and all but the largest eigenvalue of A, which is always 1, have modulus at most λ. The smallest value of λ for which this holds is denoted by $\lambda(G)$. Spectral expansion then refers to the property that $\lambda(G)$ is much smaller than 1, in which case G is referred

Tom Bannink: tombannink@gmail.com
Jop Briët: j.briet@cwi.nl
Farrokh Labib: labib@cwi.nl
Hans Maassen: H.Maassen@math.ru.nl

