

arXiv:1909.02383v3 [quant-ph] 20 Feb 2023

Quantum Brascamp-Lieb Dualities

Mario Berta,[1] David Sutter,[2] and Michael Walter[3,4]

[1]*Institute for Quantum Information, RWTH Aachen University, Aachen, Germany*
[2]*IBM Quantum, IBM Research Europe, Zürich, Switzerland*
[3]*Faculty of Computer Science, Ruhr University Bochum, Germany*
[4]*Korteweg-de Vries Institute for Mathematics, Institute for Theoretical Physics, Institute for Language, Logic & Computation, and QuSoft, University of Amsterdam, Netherlands*

Brascamp-Lieb inequalities are entropy inequalities which have a dual formulation as generalized Young inequalities. In this work, we introduce a fully quantum version of this duality, relating quantum relative entropy inequalities to matrix exponential inequalities of Young type. We demonstrate this novel duality by means of examples from quantum information theory — including entropic uncertainty relations, strong data-processing inequalities, super-additivity inequalities, and many more. As an application we find novel uncertainty relations for Gaussian quantum operations that can be interpreted as quantum duals of the well-known family of 'geometric' Brascamp-Lieb inequalities.

I. INTRODUCTION

The classical Brascamp-Lieb (BL) problem asks, given a finite sequence of surjective linear maps $L_k\colon \mathbb{R}^m \to \mathbb{R}^{m_k}$ and $q_k \in \mathbb{R}_+$ for $k \in [n]$, for the optimal constant $C \in \mathbb{R}$ such that [7, 10, 14, 47]

$$\int_{\mathbb{R}^m} \prod_{k=1}^n f_k\big(L_k x\big)\,\mathrm{d}x \leq \exp(C) \prod_{k=1}^n \|f_k\|_{1/q_k} \tag{1}$$

holds for all non-negative functions $f_k\colon \mathbb{R}^{m_k} \to \mathbb{R}_+$, $k \in [n]$, where $\|\cdot\|_p$ denotes the p-norm. Many classical integral inequalities fall into this framework, such as the Hölder inequality, Young's inequality, and the Loomis-Whitney inequality. A celebrated theorem by Lieb asserts that the optimal constant in Eq. (1) can be computed by optimizing over centred Gaussians f_k alone [47].

Remarkably, Eq. (1) has a dual, entropic formulation in terms of the *differential entropy* $H(g) := -\int g(x)\log g(x)\,\mathrm{d}x$. Namely, Eq. (1) holds for all $f_1, \ldots, f_n$ as above if, and only if, for all probability densities g on $\mathbb{R}^m$ with finite differential entropy, we have [19]

$$H(g) \leq \sum_{k=1}^n q_k H(g_k) + C\,. \tag{2}$$

Here, g_k denotes the marginal probability density on $\mathbb{R}^{m_k}$ corresponding to L_k, i.e., the push-forward of g along L_k defined by $\int_{\mathbb{R}^m} \phi(L_k x) g(x) \mathrm{d}x = \int_{\mathbb{R}^{m_k}} \phi(y) g_k(y) \mathrm{d}y$ for all bounded, continuous functions ϕ on $\mathbb{R}^{m_k}$. The duality between Eq. (1) and Eq. (2) readily generalizes to arbitrary measure spaces and measurable maps [19].

Of particular interest is the so-called *geometric* case where each L_k is a surjective partial isometry and $\sum_{k=1}^n q_k\, L_k^\dagger L_k = \mathbb{1}_{\mathbb{R}^m}$ [2–7]. In this case, Eq. (1) and Eq. (2) hold with $C = 0$. This setup includes the Hölder and Loomis-Whitney inequalities. Equivalently, we are given n subspaces $V_k \subseteq \mathbb{R}^m$ (the supports of the L_k) such that $\sum_{k=1}^n q_k\, \Pi_k = \mathbb{1}_{\mathbb{R}^m}$, where Π_k denotes

