

2022–2024

Fundamental limitations to key distillation from Gaussian states with Gaussian operations

Ludovico Lami [1,2,3,*], Ladislav Mišta, Jr. [4,†] and Gerardo Adesso [1,‡]

[1] *School of Mathematical Sciences and Centre for the Mathematics and Theoretical Physics of Quantum Non-Equilibrium Systems (CQNE), University of Nottingham, University Park, Nottingham NG7 2RD, United Kingdom*

[2] *Institut für Theoretische Physik und IQST, Universität Ulm, Albert-Einstein-Allee 11, D-89069 Ulm, Germany*

[3] *QuSoft, Korteweg-de Vries Institute for Mathematics, and Institute for Theoretical Physics, University of Amsterdam, Science Park, 1098 XG Amsterdam, the Netherlands*

[4] *Department of Optics, Palacký University, 17. listopadu 12, 771 46 Olomouc, Czech Republic*

We establish fundamental upper bounds on the amount of secret key that can be extracted from quantum Gaussian states by using only local Gaussian operations, local classical processing, and public communication. For one-way public communication, or when two-way public communication is allowed but Alice and Bob first perform destructive local Gaussian measurements, we prove that the key is bounded by the Rényi-2 Gaussian entanglement of formation $E_{F,2}^{\mathrm{G}}$. Since the inequality is saturated for pure Gaussian states, this yields an operational interpretation of the Rényi-2 entropy of entanglement as the secret key rate of pure Gaussian states that is accessible with Gaussian operations and one-way communication. In the general setting of two-way communication and arbitrary interactive protocols, we argue that $2E_{F,2}^{\mathrm{G}}$ is still an upper bound on the extractable key. We conjecture that the factor of 2 is spurious, which would imply that $E_{F,2}^{\mathrm{G}}$ coincides with the secret key rate of Gaussian states under Gaussian measurements and two-way public communication. We use these results to prove a gap between the secret key rates obtainable with arbitrary versus Gaussian operations. Such a gap is observed for states produced by sending one half of a two-mode squeezed vacuum through a pure loss channel, in the regime of sufficiently low squeezing or sufficiently high transmissivity. Finally, for a wide class of Gaussian states that includes all two-mode states, we prove a recently proposed conjecture on the equality between $E_{F,2}^{\mathrm{G}}$ and the Gaussian intrinsic entanglement. The unified entanglement quantifier emerging from such an equality is then endowed with a direct operational interpretation as the value of a quantum teleportation game.

arXiv:2010.15729v2 [quant-ph] 30 Sep 2023

I. INTRODUCTION

Quantum entanglement enables distant parties to generate a shared secret key by employing public discussion only [1–3], a feat impossible in the classical setting [4, 5] without additional assumptions on the information available to the eavesdropper [5–9]. In the last decades, quantum key distribution (QKD) has established itself as a fundamental primitive in quantum cryptography, thus gaining a central role in the flourishing quantum information science and technology [10]. Accordingly, the amount of secret key that can be extracted from a state is regarded as an entanglement measure of fundamental operational importance [11–14].

Continuous variable (CV) platforms, based on communication over quantum optical modes [15, 16], transmitted either via optical fibres or across free space [17, 18], have been of paramount importance in the demonstration of QKD. Recently, they witnessed impressive experimental progress [19–22] and will likely play a major role in any future large-scale technological implementation of QKD [23–25].

Paradigmatic examples of CV QKD protocols are those based on Gaussian states and Gaussian measurements [26–37]. The main advantage of this all-Gaussian paradigm [23, 38, 39] is that it is relatively experimentally friendly: coherent states [40–43], squeezed states [44–49], homodyne and heterodyne detection [15, 39] are nowadays relatively inexpensive ingredients, especially compared to general quantum states and operations. At the same time, it is still quite powerful in the context of QKD: in fact, it has been shown that any sufficiently entangled Gaussian state can be used, in combination with local Gaussian operations and public communication, to distil a secret key [50–52]. The effectiveness of the all-Gaussian paradigm in QKD is in stark contrast with its fundamentally limited performances at many other important tasks, such as universal quantum computation [53–56], entanglement distillation [57–59], error correction [60], and state transformations in general resource theories [61, 62].

In this paper, we investigate the operational effectiveness of the all-Gaussian framework in the context of QKD, establishing ultimate limitations on the amount of secret key that can be extracted from arbitrary multi-mode Gaussian states by means of local Gaussian operations, local classical processing, and public communication — a quantity that we call *Gaussian secret key*. The fact that the initial state is Gaussian and that the available *quantum* operations are Gaussian does not mean that the state will be Gaussian at all stages of the protocol, essentially because local classical operations are entirely unrestricted. For example, Alice could decide to apply a random displacement to her system (say, either $+s$ or $-s$, with equal probabilities), making the resulting state non-Gaussian.

In a nutshell we prove that, while key distillation is indeed possible in the Gaussian setting, it is not as efficient as it could be if also non-Gaussian measurements were allowed. Our bounds are given in terms of a Gaussian entanglement measure known as the *Rényi-2 Gaussian entanglement of formation* (denoted $E_{F,2}^{\mathrm{G}}$) [63–68], and thus endow this quantity with a sound operational meaning. In this context, after formalizing basic definitions on CV systems (Section II) and Gaussian key distillation protocols (Section III), we establish three main

* ludovico.lami@gmail.com
† mista@optics.upol.cz
‡ gerardo.adesso@nottingham.ac.uk