



Continuous Bose-Einstein condensation

Chun-Chia Chen (陳俊嘉),[1] Rodrigo González Escudero,[1] Jiří Minář,[2,3] Benjamin Pasquiou,[1,3] Shayne Bennetts,[1,3] and Florian Schreck[1,3,*]

[1] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098XH Amsterdam, The Netherlands*
[2] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098XH Amsterdam, The Netherlands*
[3] *QuSoft, Science Park 123, 1098XG Amsterdam, The Netherlands*

(Dated: September 17, 2021)

Bose-Einstein condensates (BECs) are macroscopic coherent matter waves that have revolutionized quantum science and atomic physics. They are essential to quantum simulation [1] and sensing [2, 3], for example underlying atom interferometers in space [4] and ambitious tests of Einstein's equivalence principle [5, 6]. The key to dramatically increasing the bandwidth and precision of such matter-wave sensors lies in sustaining a coherent matter wave indefinitely. Here we demonstrate continuous Bose-Einstein condensation by creating a continuous-wave (CW) condensate of strontium atoms that lasts indefinitely. The coherent matter wave is sustained by amplification through Bose-stimulated gain of atoms from a thermal bath. By steadily replenishing this bath while achieving 1000x higher phase-space densities than previous works [7, 8], we maintain the conditions for condensation. This advance overcomes a fundamental limitation of all atomic quantum gas experiments to date: the need to execute several cooling stages time-sequentially. Continuous matter-wave amplification will make possible CW atom lasers, atomic counterparts of CW optical lasers that have become ubiquitous in technology and society. The coherence of such atom lasers will no longer be fundamentally limited by the atom number in a BEC and can ultimately reach the standard quantum limit [9–11]. Our development provides a new, hitherto missing piece of atom optics, enabling the construction of continuous coherent matter-wave devices. From infrasound gravitational wave detectors [12, 13] to optical clocks [14, 15], the dramatic improvement in coherence, bandwidth and precision now within reach will be decisive in the creation of a new class of quantum sensors.

arXiv:2012.07605v2 [cond-mat.quant-gas] 16 Sep 2021

Interferometers listen to gravitational waves, image the shadow of black holes, and are at the core of countless other high-precision measurements in science and technology. An interferometer achieves its outstanding sensitivity through the interference of waves. For example, the wave-like properties of matter are exploited in the world's most accurate sensors of acceleration and gravity [16]. Interferometers reach optimal sensitivity and bandwidth by using continuous, coherent, high-brightness sources, which is why many optical interferometers, such as LIGO, are based on CW lasers. However, no continuous matter-wave source with long-range coherence currently exists, which prevents atom interferometers from reaching their full potential. Creating such a CW source of matter will benefit applications ranging from geodesy [17, 18], to tests of Einstein's equivalence principle [5, 6], gravitational-wave detection [12, 13, 19–21], and dark-matter and dark-energy searches [22, 23].

The key to realizing a CW BEC of atoms is to continuously amplify the atomic matter wave while preserving its phase coherence [24]. An amplification process is essential to compensate naturally occurring atom losses, e.g. from molecule formation. It is also needed to replace the atoms that will be coupled out of the BEC for sustaining an atom laser or atom interferometer. Addressing this challenge requires two ingredients: a gain mechanism that amplifies the BEC and a continuous supply of ultracold atoms near quantum degeneracy.

The first steps towards a continuous gain mechanism were taken by [25], where merging of independent condensates periodically added atoms to an already existing BEC, but where coherence was not retained across merger events. A Bose-stimulated gain mechanism into a single dominant mode (the BEC) is required to provide gain without sacrificing phase coherence. Such gain mechanisms have been demonstrated using elastic collisions between thermal atoms [24, 26], stimulated photon emission [27], four wave mixing [28, 29] or superradiance [30]. However, in all these demonstrations the gain mechanism could not be sustained indefinitely.

To sustain gain the second ingredient is needed: a continuous supply of ultracold, dense gas with a phase-space density — the occupancy of the lowest motional quantum state — approaching $\rho = 1$. Great efforts were spent developing continuously cooled beams of atoms [31, 32] and continuously loaded traps [7, 8], which so far have reached phase-space densities of $\rho = 10^{-3}$. To achieve the required µK temperatures, these experiments have to use laser cooling, but near-resonant laser cooling light is highly destructive for BECs [33]. Several experiments have maintained a BEC in the presence of harmful light, either by spatially separating the laser cooling from the quantum gas [7, 25, 31, 32, 34] or by reducing the quantum gas' absorbance [24, 35–37].

Here we demonstrate the creation of a CW BEC that

* ContinuousBEC@strontiumBEC.com