

Holographic entanglement negativity and replica symmetry breaking

Xi Dong,[a] **Xiao-Liang Qi,**[b] **and Michael Walter**[c]

[a] *Department of Physics, University of California, Santa Barbara, CA 93106, USA*

[b] *Stanford Institute for Theoretical Physics, Physics Department, Stanford University, Stanford, CA 94305, USA*

[c] *Korteweg-de Vries Institute for Mathematics, Institute for Theoretical Physics, Institute for Logic, Language, and Computation, QuSoft, University of Amsterdam, The Netherlands*

E-mail: xidong@ucsb.edu, xlqi@stanford.edu, m.walter@uva.nl

arXiv:2101.11029v2 [hep-th] 7 Jun 2021

Abstract: Since the work of Ryu and Takayanagi, deep connections between quantum entanglement and spacetime geometry have been revealed. The negative eigenvalues of the partial transpose of a bipartite density operator is a useful diagnostic of entanglement. In this paper, we discuss the properties of the associated *entanglement negativity* and its Rényi generalizations in holographic duality. We first review the definition of the Rényi negativities, which contain the familiar logarithmic negativity as a special case. We then study these quantities in the random tensor network model and rigorously derive their large bond dimension asymptotics. Finally, we study entanglement negativity in holographic theories with a gravity dual, where we find that Rényi negativities are often dominated by bulk solutions that break the replica symmetry. From these replica symmetry breaking solutions, we derive general expressions for Rényi negativities and their special limits including the logarithmic negativity. In fixed-area states, these general expressions simplify dramatically and agree precisely with our results in the random tensor network model. This provides a concrete setting for further studying the implications of replica symmetry breaking in holography.

