

2021–2022

arXiv:2101.11032v5 [quant-ph] 20 Oct 2021

The original Wigner's friend paradox within a realist toy model

Matteo Lostaglio[1] and Joseph Bowles[2]

[1] *Korteweg-de Vries Institute for Mathematics and QuSoft, University of Amsterdam, The Netherlands*
[2] *ICFO - Institut de Ciències Fotòniques, The Barcelona Institute of Science and Technology, 08860 Castelldefels, Spain*

The original Wigner's friend paradox is a gedankenexperiment involving an observer described by an external agent. The paradox highlights the tension between unitary evolution and collapse in quantum theory, and is sometimes taken as requiring a reassessment of the notion of objective reality. In this note however we present a classical toy model in which (i) The contradicting predictions at the heart of the thought experiment are reproduced (ii) Every system is in a well-defined state at all times. The toy model shows how puzzles such as Wigner's friend's experience of being in a superposition, conflicts between different agents' descriptions of the experiment, the positioning of the Heisenberg's cut and the apparent lack of objectivity of measurement outcomes can be explained within a classical model where there exists an objective state of affairs about every physical system at all times. Within the model, the debate surrounding the original Wigner's friend thought experiment and its resolution have striking similarities with arguments concerning the nature of the second law of thermodynamics. The same conclusion however does not apply to more recent extensions of the gedankenexperiment featuring multiple encapsulated observers, and shows that such extensions are indeed necessary avoid simple classical explanations.

I. SETTING THE STAGE

The original Wigner's friend gedankenexperiment is a thought experiment first considered by Wigner in 1961 [1] that highlights the tension between unitary evolution and measurement collapse in quantum theory. It involves two agents, namely Wigner (W) and his friend (F).

In the simplest setting, a single qubit system is initially prepared in the state $|0\rangle_S$. S undergoes a (Hadamard) unitary evolution, after which its state is $\frac{1}{\sqrt{2}}(|0\rangle_S+|1\rangle_S)$. At this point S is measured by Wigner's friend, who finds it in state $|0\rangle_S$ or $|1\rangle_S$ with equal probability according to the Born rule.

Wigner is a superobserver sitting outside his friend's lab. According to him, the evolution of both S and his friend F can be described by Schrödinger's equation. In the most bare-bone description, W describes the relevant degrees of freedom of F as being themselves a two-level system, schematically representing a memory on which the measurement outcome is imprinted. The friend's memory state is initialised in $|0\rangle_F$ and the measurement of S by F can be described as a CNOT between the two:

$$\frac{1}{\sqrt{2}}(|0\rangle_S+|1\rangle_S)\otimes|0\rangle_F \mapsto \frac{1}{\sqrt{2}}(|0\rangle_S\otimes|0\rangle_F+|1\rangle_S\otimes|1\rangle_F) := |\phi^+\rangle_{SF}.$$

After the interaction, the states $|0\rangle_F$ and $|1\rangle_F$ correspond to the labels "the friend sees outcome 0" and "the friend sees outcome 1". Wigner can verify his prediction by performing a measurement in the basis of 2-qubit maximally entangled states $\{|\psi^\pm\rangle_{SF}, |\phi^\pm\rangle_{SF}\}$. He will get the outcome ϕ^+ with probability 1. Hence, if quantum theory is correct at all scales, the thought experiment requires that Wigner's description of the global entangled state must be compatible with Wigner's friend experience of a definite measurement outcome.

Note however that if F applies the collapse rule she would assign to S the state $|0\rangle_S$ *or* $|1\rangle_S$. According to this procedure she would then predict that the two outcomes $|\phi^\pm\rangle_{SF}$ of Wigner's measurement can occur, which is in contradiction with W's prediction of obtaining ϕ^+ with probability 1. Since we assume here and throughout the universality of quantum mechanics, this inference from the collapse rule must be unwarranted.

Available explanations of this thought experiment compatible with quantum mechanics involve serious departures from the classical worldview. Broadly speaking they are of two kinds: perspectival interpretations such as QBism [2], which are at ease with the idea that every fact, including a measurement outcome, is relative to a particular observer; and interpretations such as Bohmian mechanics [3] or the many-world interpretation [4], in which the quantum state is taken to be part of a highly non-classical ontology and there is no actual collapse, which is understood as an effective procedure. So it is natural to ask whether classical models reproducing the same phenomenology exist, and how the abovementioned issues look like within such descriptions.

The plan

The aim of this note is two-fold:

1. Highlight that the original Wigner's friend paradox admits a classical explanation. That is, the contradicting predictions at the core of the gedankenexperiment arise in a simple non-contextual classical model, in which different agents have different (and conflicting) descriptions. Here, 'contradicting predictions' refers to the fact that Wigner's and the friend's descriptions give different predictions for the possible outcomes of Wigner's entangled measurement. While this will not come as a surprise in the quantum foundations community, we think it is worth to present it in some detail, since it shows that recent 'supercharged' extensions of Wigner's friend [5–9] are necessary to avoid simple classical explanations.

