

arXiv:2102.01064v1 [quant-ph] 1 Feb 2021

Quantum Gravity in the Lab: Teleportation by Size and Traversable Wormholes, Part II

Sepehr Nezami,[1,2,3] Henry W. Lin,[2,4] Adam R. Brown,[2,3,5] Hrant Gharibyan,[1] Stefan Leichenauer,[2,5] Grant Salton,[6,7,1,3] Leonard Susskind,[3,2,5] Brian Swingle,[8] and Michael Walter[9]

[1] *Institute for Quantum Information and Matter, Caltech, Pasadena, CA 91125, USA*
[2] *Google, Mountain View, CA 94043, USA*
[3] *Department of Physics, Stanford University, Stanford, CA 94305, USA*
[4] *Physics Department, Princeton University, Princeton, NJ 08540, USA*
[5] *Sandbox@Alphabet, Mountain View, CA 94043, USA*
[6] *Amazon Quantum Solutions Lab, Seattle, WA 98170, USA*
[7] *AWS Center for Quantum Computing, Pasadena, CA 91125, USA*
[8] *Brandeis University, Waltham, MA 02453, USA,*
University of Maryland, College Park, MD 20742, USA
[9] *Korteweg-de Vries Institute for Mathematics, Institute for Theoretical Physics, Institute for Logic, Language and Computation & QuSoft, University of Amsterdam, The Netherlands*

In [1] we discussed how quantum gravity may be simulated using quantum devices and gave a specific proposal—teleportation by size and the phenomenon of size-winding. Here we elaborate on what it means to do 'Quantum Gravity in the Lab' and how size-winding connects to bulk gravitational physics and traversable wormholes. Perfect size-winding is a remarkable, fine-grained property of the size wavefunction of an operator; we show from a bulk calculation that this property must hold for quantum systems with a nearly-AdS_2 bulk. We then examine in detail teleportation by size in three systems: the Sachdev-Ye-Kitaev model, random matrices, and spin chains, and discuss prospects for realizing these phenomena in near-term quantum devices.

