

2021–2022

arXiv:2111.12130v2 [quant-ph] 5 Aug 2022

Continuous thermomajorization and a complete set of laws for Markovian thermal processes

Matteo Lostaglio*,[1,2] and Kamil Korzekwa*,[3]

[1] *Korteweg-de Vries Institute for Mathematics and QuSoft, University of Amsterdam, The Netherlands*
[2] *QuTech, Delft University of Technology, P.O. Box 5046, 2600 GA Delft, The Netherlands*
[3] *Faculty of Physics, Astronomy and Applied Computer Science, Jagiellonian University, 30-348 Kraków, Poland*

The standard dynamical approach to quantum thermodynamics is based on Markovian master equations describing the thermalization of a system weakly coupled to a large environment, and on tools such as entropy production relations. Here we develop a new framework overcoming the limitations that the current dynamical and information theory approaches encounter when applied to this setting. More precisely, we introduce the notion of continuous thermomajorization, and employ it to obtain necessary and sufficient conditions for the existence of a Markovian thermal process transforming between given initial and final energy distributions of the system. These lead to a complete set of generalized entropy production inequalities including the standard one as a special case. Importantly, these conditions can be reduced to a finitely verifiable set of constraints governing non-equilibrium transformations under master equations. What is more, the framework is also constructive, i.e., it returns explicit protocols realizing any allowed transformation. These protocols use as building blocks elementary thermalizations, which we prove to be universal controls. Finally, we also present an algorithm constructing the full set of energy distributions achievable from a given initial state via Markovian thermal processes and provide a `Mathematica` implementation solving $d = 6$ on a laptop computer in minutes.

I. INTRODUCTION

It is well-known that thermodynamics can be formulated in the resource theory language of information theory [1–6]. Since it focuses only on input-output relations under a class of quantum operations, the resource theory is unable to discuss how the process is realized in time, nor does it involve the notions of entropy production or master equations, even though the latter are commonplace in standard approaches. As such, the standard and the resource-theoretic frameworks are for the most part disconnected. Somewhat surprisingly, despite all its powerful theorems, the resource theory has not affected the analysis of a thermodynamics practitioner following the more explicit dynamical approaches based on the master equation formalism [7].

The aim of this work is to overcome this situation by showing how to unify the master equations and information theory tools. We first summarize the basic notions of both approaches to thermodynamics in Sec. II, and demonstrate the insufficiency of each of them, when taken separately, to capture relevant thermodynamic constraints. Then, in Sec. III, we propose a hybrid approach that can overcome these limitations. In constructing our solution we highlight the necessity of finding a *finitely verifiable* set of thermodynamic laws specifying when one state can be thermodynamically transformed into another one. Furthermore, we wish to go beyond the question of *whether* a thermodynamic transformation exists, and ask *how* it should be realized. The resource theory approaches are for the most part silent about the latter, which is arguably a central obstacle to their application to concrete problems. Our framework, on the contrary, will be *constructive*.

In Sec. IV we introduce our main novel technical tool, *continuous thermomajorization*, which extends the concept of thermomajorization introduced in Refs. [3, 8]. We prove that this partial order of energy distributions provides necessary and sufficient conditions for the existence of a thermalization process generated by a Markovian master equation. More precisely, we show that an out of equilibrium energy distribution can be transformed into another one by a Markovian thermal process if and only if the former continuously thermomajorizes the latter. We then connect with the notion of entropy production [9] in Sec. V, where we show that continuous thermomajorization allows one to identify a *complete* set of generalized entropy production relations, including the standard one as a special case.

Our ultimate goal, however, is to find a finitely-verifiable set of such conditions. To achieve this, we first prove in Sec. VI that *elementary thermalizations*, generated by simple reset master equations on two-level submanifolds, form universal controls in the Markovian regime. In other words, every state transformation that can be achieved by a Markovian thermal process can also be achieved by a sequence of elementary thermalizations. Employing this simplified set of controls, in Sec. VII we show that continuous thermomajorization indeed can be checked in a finite number of steps. Remarkably, we can also return the exact sequence of elementary controls required to realize any allowed transformation. As a final result of our paper, in Sec. VIII we provide an explicit



* These authors contributed equally to this work.
Emails: lostaglio@protonmail.com, korzekwa.kamil@gmail.com