



Narrow-line imaging of single strontium atoms in shallow optical tweezers

Alexander Urech,[1,2] Ivo H. A. Knottnerus,[1,2,3] Robert J. C. Spreeuw,[1,2] and Florian Schreck[1,2,*]

[1] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098XH Amsterdam, The Netherlands*
[2] *QuSoft, Science Park 123, 1098XG Amsterdam, The Netherlands*
[3] *Eindhoven University of Technology, P.O. Box 513, 5600MB Eindhoven, The Netherlands*

(Dated: February 14, 2022)

Single strontium atoms held in optical tweezers have so far only been imaged using the broad $^1S_0$ - $^1P_1$ transition. For Yb, use of the narrow (183 kHz-wide) $^1S_0$ - $^3P_1$ transition for simultaneous imaging and cooling has been demonstrated in tweezers with a magic wavelength for the imaging transition. We demonstrate high-fidelity imaging of single Sr atoms using its even narrower (7.4 kHz-wide) $^1S_0$ - $^3P_1$ transition. The atoms are trapped in *non*-magic-wavelength tweezers. We detect the photons scattered during Sisyphus cooling, thus keeping the atoms near the motional ground state of the tweezer throughout imaging. The fidelity of detection is 0.9991(4) with a survival probability of 0.97(2). An atom in a tweezer can be held under imaging conditions for 79(3) seconds allowing for hundreds of images to be taken, limited mainly by background gas collisions. We detect atoms in an arrary of 36 tweezers with 813.4-nm light and trap depths of 135(20) μK. This trap depth is three times shallower than typically used for imaging on the broad $^1S_0$ - $^1P_1$ transition. Narrow-line imaging opens the possibility to even further reduce this trap depth, as long as all trap frequencies are kept larger than the imaging transition linewidth. Imaging using a narrow-linewidth transition in a non-magic-wavelength tweezer also allows for selective imaging of a given tweezer. As a demonstration, we selectively image (hide) a single tweezer from the array. This provides a useful tool for quantum error correction protocols.

arXiv:2202.05727v1 [physics.atom-ph] 11 Feb 2022

I. INTRODUCTION

Optical tweezers have emerged as a powerful tool for quantum applications. They enable state of the art quantum simulation and computation [1–4], high fidelity and long coherence time qubits [5–8], quantum metrology [9–11], quantum chemistry [12, 13], among numerous other applications. Optical tweezers with alkaline-earth(-like) atoms, in particular with strontium and ytterbium, have been recently realized, offering new possibilities in expanding these applications [14–16].

In all strontium tweezer experiments demonstrated so far, the fluorescence of single atoms on the broad (30 MHz) $^1S_0$ - $^1P_1$ transition at 461 nm was recorded, while simultaneously cooling the atoms on the narrow (7.4 kHz) $^1S_0$ - $^3P_1$ transition at 689 nm [4–6, 9–11, 14, 15, 17]. This 'blue imaging' method allows for high-fidelity detection of single atoms in tweezers with high survival probability [14, 15, 17]. However, blue imaging requires repumpers to close the 5s4d $^1D_2$ decay channel, which can only be done at tweezer wavelengths where also the $^1D_2$ state is trapped [5, 6, 9, 17]. Furthermore, the slightly higher scattering rate obtained in the blue imaging process ($\sim$ 75 kHz) can only be used as long as the tweezers are sufficiently deep. Any excess heating from the imaging process can then be cooled away after the image. As the trap depth of the tweezer is reduced, the advantage of fast imaging is lost because the scattering rate must also be reduced to balance heating from imaging and cooling. Reducing the tweezer trap depth has the advantages of decreased laser power requirement per tweezer (allowing for more tweezers using a given laser source) and increasing metastable state lifetimes (reduced off-resonant scattering of tweezer light).

A simpler method for imaging alkaline-earth(-like) atoms in tweezers is to use the narrow $^1S_0$ - $^3P_1$ transition for both cooling and imaging. Single atom detection by fluorescence imaging on a (less) narrow transition has previously been demonstrated in ytterbium for two different isotopes. In both cases tweezers with a magic-wavelength for the imaging transition were used [7, 16].

Here we detect single ^{88}Sr atoms using only the $^1S_0$ - $^3P_1$ transition for simultaneous'red imaging' and cooling. We use optical tweezers that are *non*-magic for the imaging transition, but magic for the Sr clock transition ($^1S_0$ - $^3P_0$). We detect the photons scattered during an *attractive* Sisyphus cooling process [17], thus keeping the atoms near the motional ground state of the tweezer throughout imaging.

Attractive Sisyphus cooling is possible at tweezer wavelengths where the excited state experiences a deeper trap depth than the ground state. More specifically, at our tweezer wavelength of 813.4 nm, the excited state ($|e\rangle \equiv {}^3P_1\,(|m_j| = 1)$) confinement is 1.24 times greater than the ground state ($|g\rangle \equiv {}^1S_0$), see Fig. 1(a). This cooling process can be very efficient with a proper choice of parameters, leading to a large reduction in energy per scattered photon, and a small number of scattered photons needed to cool the atom [18]. In addition, due to the anharmonicity of the tweezer potential, atomic motion in all directions is coupled, allowing for a single radial cooling beam to remove energy from all directions [17, 18].

With balanced heating and cooling from the imaging

* NarrowLineSrTweezerImaging@strontiumBEC.com