

2022–2024

arXiv:2204.06800v2 [quant-ph] 22 Jul 2022

$9 \times 4 = 6 \times 6$: Understanding the quantum solution to Euler's problem of 36 officers

Karol Życzkowski[1,2], **Wojciech Bruzda**[1], **Grzegorz Rajchel-Mieldzioć**[3,2], **Adam Burchardt**[4,1], **Suhail Ahmad Rather**[5], **Arul Lakshminarayan**[5]

[1] Institute of Theoretical Physics, Jagiellonian University, ul. Łojasiewicza 11, 30-348 Kraków, Poland
[2] Center for Theoretical Physics, Polish Academy of Sciences, 02-668 Warszawa, Poland
[3] ICFO-Institut de Ciencies Fotoniques, The Barcelona Institute of Science and Technology, Av. Carl Friedrich Gauss 3, 08860 Castelldefels (Barcelona), Spain
[4] QuSoft, CWI and University of Amsterdam, Science Park 123, 1098 XG Amsterdam, the Netherlands
[5] Department of Physics, Indian Institute of Technology Madras, Chennai 600036, India

Abstract. The famous combinatorial problem of Euler concerns an arrangement of 36 officers from six different regiments in a 6×6 square array. Each regiment consists of six officers each belonging to one of six ranks. The problem, originating from Saint Petersburg, requires that each row and each column of the array contains only one officer of a given rank and given regiment. Euler observed that such a configuration does not exist. In recent work, we constructed a solution to a quantum version of this problem assuming that the officers correspond to quantum states and can be entangled. In this paper, we explain the solution which is based on a partition of 36 officers into nine groups, each with four elements. The corresponding quantum states are locally equivalent to maximally entangled two-qubit states, hence each officer is entangled with at most three out of his 35 colleagues. The entire quantum combinatorial design involves 9 Bell bases in nine complementary 4–dimensional subspaces.

1. Introduction

Designs composed of elements of a finite set arranged with certain symmetry and balance have been known since antiquity. Nowadays, they are commonly referred to as *combinatorial designs* [1], mostly developed in the 18th century along with the general growth of combinatorics. Some of them, such as magic squares and Sudoku grids, offer mathematical recreations. However, the 20th century brought to light several applications of combinatorics to real-world problems. The theory of combinatorial designs

