

2021–2022

arXiv:2206.11783v3 [quant-ph] 29 Sep 2023

Kirkwood-Dirac quasiprobability approach to the statistics of incompatible observables

Matteo Lostaglio[1], Alessio Belenchia[2,3], Amikam Levy[4], Santiago Hernández-Gómez[5,6,7], Nicole Fabbri[5,7], and Stefano Gherardini[8,5]

[1]Korteweg-de Vries Institute for Mathematics and QuSoft, University of Amsterdam, The Netherlands
[2]Institut für Theoretische Physik, Eberhard-Karls-Universität Tübingen, 72076 Tübingen, Germany
[3]Centre for Theoretical Atomic, Molecular and Optical Physics, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, United Kingdom
[4]Department of Chemistry and Center for Quantum Entanglement Science and Technology, Bar-Ilan University, Ramat-Gan 52900, Israel
[5]European Laboratory for Non-linear Spectroscopy (LENS), Università di Firenze, I-50019 Sesto Fiorentino, Italy
[6]Dipartimento di Fisica e Astronomia, Università di Firenze, I-50019, Sesto Fiorentino, Italy
[7]Istituto Nazionale di Ottica del Consiglio Nazionale delle Ricerche (CNR-INO), I-50019 Sesto Fiorentino, Italy
[8]Istituto Nazionale di Ottica del Consiglio Nazionale delle Ricerche (CNR-INO), Area Science Park, Basovizza, I-34149 Trieste, Italy

Recent work has revealed the central role played by the Kirkwood-Dirac quasiprobability (KDQ) as a tool to properly account for non-classical features in the context of condensed matter physics (scrambling, dynamical phase transitions) metrology (standard and post-selected), thermodynamics (power output and fluctuation theorems), foundations (contextuality, anomalous weak values) and more. Given the growing relevance of the KDQ across the quantum sciences, our aim is two-fold: First, we highlight the role played by quasiprobabilities in characterizing the statistics of quantum observables and processes in the presence of measurement incompatibility. In this way, we show how the KDQ naturally underpins and unifies quantum correlators, quantum currents, Loschmidt echoes, and weak values. Second, we provide novel theoretical and experimental perspectives by discussing a wide variety of schemes to access the KDQ and its non-classicality features.

Matteo Lostaglio: lostaglio@protonmail.com, These authors contributed equally to this work

Alessio Belenchia: These authors contributed equally to this work

Contents

1 Introduction **2**

2 Why quasiprobabilities? **3**
2.1 A no-go theorem on the characterization of incompatible observables in quantum theory 3
2.2 The no-go theorem in a thermodynamic context 4

3 The physics behind the Kirkwood-Dirac quasiprobability **5**
3.1 The quasiprobability behind quantum correlators 5
3.2 Linear response theory 6
3.3 Quantum currents 8
3.4 Loschmidt echo 8
3.5 Weak values are conditional Kirkwood-Dirac quasiprobability averages 9
3.6 Non-commutatitivity in thermodynamics and the Kirkwood-Dirac quasiprobability 9

4 Measuring the Kirkwood-Dirac quasiprobability **10**
4.1 Weak two-point-measurement protocol 11

