

arXiv:2209.13209v1 [physics.atom-ph] 27 Sep 2022

Tracking the Vector Acceleration with a Hybrid Quantum Accelerometer Triad

Simon Templier,[1,2] Pierrick Cheiney,[1] Quentin d'Armagnac de Castanet,[1,2] Baptiste Gouraud,[1] Henri Porte,[1] Fabien Napolitano,[1] Philippe Bouyer,[1,3,4,5] Baptiste Battelier,[2,*] and Brynle Barrett[1,2,5]

[1]iXblue, 34 rue de la Croix de Fer, 78105 Saint-Germain-en-Laye, France

[2]LP2N, Laboratoire Photonique, Numérique et Nanosciences, Université Bordeaux–IOGS–CNRS:UMR 5298, 1 rue François Mitterrand, 33400 Talence, France

[3]Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, Science Park 904, 1098XH Amsterdam, The Netherlands

[4]QuSoft, Science Park 123, 1098XG Amsterdam, The Netherlands

[5]Eindhoven University of Technology, The Netherlands

[5]Department of Physics, University of New Brunswick, 8 Bailey Dr., Fredericton NB, E3B 5A3, Canada

*To whom correspondence should be addressed; E-mail: baptiste.battelier@institutoptique.fr.

September 28, 2022

Robust and accurate acceleration tracking remains a challenge in many fields. For geophysics and economic geology, precise gravity mapping requires on-board sensors combined with accurate positioning and navigation systems. Cold-atom-based quantum inertial sensors can potentially provide such high-precision instruments. However, current scalar instruments require precise alignment with vector quantities. Here, we present the first hybrid three-axis accelerometer exploiting the quantum advantage to measure the full accelera-

