

2022–2025



arXiv:2211.16518v2 [quant-ph] 3 Aug 2023

Optimizing sparse fermionic Hamiltonians

Yaroslav Herasymenko[1,2], Maarten Stroeks[2,3], Jonas Helsen[1], and Barbara Terhal[2,3]

[1]QuSoft & CWI, Science Park 123 1098 XG Amsterdam, The Netherlands
[2]QuTech, Delft University of Technology, Lorentzweg 1, 2628 CJ Delft, The Netherlands
[3]EEMCS Faculty, Delft University of Technology, Van Mourik Broekmanweg 6, 2628 XE, Delft, The Netherlands

We consider the problem of approximating the ground state energy of a fermionic Hamiltonian using a Gaussian state. In sharp contrast to the dense case [1, 2], we prove that strictly q-local *sparse* fermionic Hamiltonians have a constant Gaussian approximation ratio; the result holds for any connectivity and interaction strengths. Sparsity means that each fermion participates in a bounded number of interactions, and strictly q-local means that each term involves exactly q fermionic (Majorana) operators. We extend our proof to give a constant Gaussian approximation ratio for sparse fermionic Hamiltonians with both quartic and quadratic terms. With additional work, we also prove a constant Gaussian approximation ratio for the so-called sparse SYK model with strictly 4-local interactions (sparse SYK-4 model). In each setting we show that the Gaussian state can be efficiently determined. Finally, we prove that the $O(n^{-1/2})$ Gaussian approximation ratio for the normal (dense) SYK-4 model extends to SYK-q for even $q > 4$, with an approximation ratio of $O(n^{1/2-q/4})$. Our results identify non-sparseness as the prime reason that the SYK-4 model can fail to have a constant approximation ratio [1, 2].

Contents

1 Introduction 1

2 Statement of results 2
2.1 Preliminaries 2
2.2 Sparse fermionic Hamiltonians 3
2.3 The sparse $q = 4$ SYK model 4
2.4 Higher-q SYK models 4

3 Discussion 5

4 Background on Gaussian states 6
4.1 Gaussian states 6

5 Approximation ratios for sparse fermionic Hamiltonians 7

6 Sparse Hamiltonians with terms of weight 2 and 4 8

7 The sparse SYK-4 model 10

8 Upper bound on Gaussian approximation ratio for SYK-q Hamiltonians 11
8.1 Gaussian upper bound for SYK-q models 11
8.2 Maximum eigenvalue lower bound for q-local SYK Hamiltonians 12

Acknowledgements 15

References 15

A Extensive sets of all anti-commuting terms 17

B Splitting sparse Hamiltonians into diffuse interaction sets 18

C Majorana matchings from diffuse interaction sets 19

D Matchings and Gaussian states 20

E Concentration bounds for sparse SYK-4 20

F Moment bound for dense SYK-q 26
F.1 Upper bound for moments of D_0 (diagonal-free contribution) 27
F.2 Upper bound for moments of D_1, D_2, D_3 and D_4 30

G Two-colored SYK to standard SYK 33

1 Introduction

Approximating the ground state energy of a local Hamiltonian is a central problem in both physics and computer science. In computer science it plays a key role in complexity theory [3], while in physics ground states capture the behaviour of systems at low energy. Two common families of Hamiltonians of interest are those defined on collections of qubits and those acting on fermionic degrees of freedom. Fermionic Hamiltonians model various physical systems, such as electrons in condensed matter and quantum chemistry — prime targets for quantum simulation. Fermions also define a model of quantum computation, equivalent to the one based on qubits [4]. Despite its practical and conceptual relevance, the general problem of approximating fermionic ground state energies is currently less well understood than its qubit counterpart.