

2022–2024

On tests of the quantum nature of gravitational interactions in presence of non-linear corrections to quantum mechanics

Giovanni Spaventa[1], Ludovico Lami[1,2,3,4], and Martin B. Plenio[1]

[1]Institute of Theoretical Physics and IQST, Universität Ulm, Albert-Einstein-Allee 11 D-89081, Ulm, Germany
[2]QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands
[3]Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands
[4]Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands

arXiv:2302.00365v2 [quant-ph] 23 Oct 2023

When two particles interact primarily through gravity and follow the laws of quantum mechanics, the generation of entanglement is considered a hallmark of the quantum nature of the gravitational interaction. However, we demonstrate that entanglement dynamics can also occur in the presence of a weak quantum interaction and non-linear corrections to local quantum mechanics, even if the gravitational interaction is classical or absent at short distances. This highlights the importance of going beyond entanglement detection to conclusively test the quantum character of gravity, and it requires a thorough examination of the strength of other quantum forces and potential non-linear corrections to quantum mechanics in the realm of large masses.

Introduction — Does the gravitational field require quantization and if so, how do we formulate the correct theory of quantum gravity? Despite intense research and debate, these key questions at the interface of quantum mechanics and general relativity, two theories that have revolutionised physics in the early parts of the 20th century, remain open to this day. These issues have been hotly discussed already at the 1957 Chapel Hill Conference on the Role of Gravitation in Physics [1]. At the time not only did Richard Feynman state that "we're in trouble if we believe in quantum mechanics but *don't* quantize gravitational theory"; he also supported this assessment with a *Gedankenexperiment*. He considered a massive particle to be placed in a coherent superposition of its spatial degrees of freedom by first preparing an internal spin degree-of-freedom of this particle in coherent superposition and allowing this particle to pass through a Stern-Gerlach apparatus. Then, crucially, he assumed this particle to interact purely gravitationally with yet another massive particle. Indeed, treating the gravitational interaction as either classical or quantum mechanical results in very different quantum states of the two particles and thus experimental results of subsequent measurements. The particles would emerge with correlated positions that are described either by an incoherent mixture or by a coherent superposition, respectively. The latter, in modern quantum information parlance referred to as an entangled state, let Feynman to conclude that "We would then have to analyze through the channel provided by the gravitational field itself via the quantum mechanical amplitudes. Therefore, there must be an amplitude for the gravitational field".

Hence, as discussed further in [2, 3, 4, 5] the observation of the generation of entanglement between two massive particles would certify the quantum mechanical character of their gravitational interaction. If this interaction is assumed to be mediated by a gravitational field and the local dynamics follows the laws of quantum mechanics, then this would falsify the assumption of a classical force carrier and thereby imply the non-classical nature of the gravitational field [6, 7, 8, 9, 10]. Note that in this argument the existence of a gravitational field is taken as an assumption that is not directly verified by the experiment [11].

Giovanni Spaventa: giovanni.spaventa@uni-ulm.de
Ludovico Lami: ludovico.lami@gmail.com
Martin B. Plenio: martin.plenio@uni-ulm.de

