

2022–2024

arXiv:2303.04949v2 [quant-ph] 1 Aug 2023

Pretty good measurement for bosonic Gaussian ensembles

Hemant K. Mishra,[1,2,*] Ludovico Lami,[3,4,5,6,†] Prabha Mandayam,[7,8,‡] and Mark M. Wilde[1,2,§]

[1]*School of Electrical and Computer Engineering, Cornell University, Ithaca, New York 14850, USA*
[2]*Hearne Institute for Theoretical Physics, Department of Physics and Astronomy, and Center for Computation and Technology, Louisiana State University, Baton Rouge, Louisiana 70803, USA*
[3]*Institute of Theoretical Physics and IQST, Universität Ulm, Albert-Einstein-Allee 11D-89069 Ulm, Germany*
[4]*QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[5]*Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*
[6]*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[7]*Center for Quantum Information, Communication and Computing, IIT Madras, Chennai 600036, India*
[8]*Department of Physics, Indian Institute of Technology Madras, Chennai 600036, India*

The pretty good measurement is a fundamental analytical tool in quantum information theory, giving a method for inferring the classical label that identifies a quantum state chosen probabilistically from an ensemble. Identifying and constructing the pretty good measurement for the class of bosonic Gaussian states is of immediate practical relevance in quantum information processing tasks. Holevo recently showed that the pretty good measurement for a bosonic Gaussian ensemble is a bosonic Gaussian measurement that attains the accessible information of the ensemble [*IEEE Trans. Inf. Theory, 66(9):5634–564, 2020*]. In this paper, we provide an alternate proof of Gaussianity of the pretty good measurement for a Gaussian ensemble of multimode bosonic states, with a focus on establishing an explicit and efficiently computable Gaussian description of the measurement. We also compute an explicit form of the mean square error of the pretty good measurement, which is relevant when using it for parameter estimation.

Generalizing the pretty good measurement is a quantum instrument, called the pretty good instrument. We prove that the post-measurement state of the pretty good instrument is a faithful Gaussian state if the input state is a faithful Gaussian state whose covariance matrix satisfies a certain condition. Combined with our previous finding for the pretty good measurement and provided that the same condition holds, it follows that the expected output state is a faithful Gaussian state as well. In this case, we compute an explicit Gaussian description of the post-measurement and expected output states. Our findings imply that the pretty good instrument for bosonic Gaussian ensembles is no longer merely an analytical tool, but that it can also be implemented experimentally in quantum optics laboratories.

Keywords: pretty good measurement; pretty good instrument; bosonic Gaussian ensemble; Gaussian measurement; exponential quadratic forms; mean square error.

Dedicated to Alexander S. Holevo on the occasion of his 80th birthday. Professor Holevo's numerous seminal contributions, from past [1] to present [2, 3], and his textbooks [4, 5] have served as an inspiration and source of knowledge for generations of quantum information scientists.



* hemant.mishra@cornell.edu
† ludovico.lami@gmail.com
‡ prabhamd@physics.iitm.ac.in
§ wilde@cornell.edu