

2022–2024

Orthonormal bases of extreme quantumness

Marcin Rudziński[1,2], Adam Burchardt[3], and Karol Życzkowski[1,4]

[1] Faculty of Physics, Astronomy and Applied Computer Science, Jagiellonian University, ul. Łojasiewicza 11, 30-348 Kraków, Poland

[2]Doctoral School of Exact and Natural Sciences, Jagiellonian University, ul. Łojasiewicza 11, 30-348 Kraków, Poland

[3]QuSoft, CWI and University of Amsterdam, Science Park 123, 1098 XG Amsterdam, the Netherlands

[4]Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warszawa, Poland

arXiv:2306.00532v3 [quant-ph] 17 Jan 2024

Spin anticoherent states acquired recently a lot of attention as the most "quantum" states. Some coherent and anticoherent spin states are known as optimal quantum rotosensors. In this work, we introduce a measure of quantumness for orthonormal bases of spin states, determined by the average anticoherence of individual vectors and the Wehrl entropy. In this way, we identify the most coherent and most quantum states, which lead to orthogonal measurements of extreme quantumness. Their symmetries can be revealed using the Majorana stellar representation, which provides an intuitive geometrical representation of a pure state by points on a sphere. Results obtained lead to maximally (minimally) entangled bases in the $2j+1$ dimensional symmetric subspace of the 2^{2j} dimensional space of states of multipartite systems composed of $2j$ qubits. Some bases found are isocoherent as they consist of all states of the same degree of spin-coherence.

1. Introduction

Geometric methods play an essential role while studying physical systems in classical mechanics [1,2], relativity [3], quantum mechanics [4,5], and quantum field theory [6]. The stellar representation, also called *the Majorana representation*, is one of the important geometrical representations in quantum mechanics [7]. The stellar representation presents a spin-j pure state in $N = 2j + 1$ dimensional Hilbert space as a collection of $2j$ points on a sphere. The same constellation represents a symmetric state of a system consisting of $2j$ qubits. In the case of a spin-$\frac{1}{2}$ particle, it reduces to the celebrated Bloch representation of a two-level quantum system (qubit). This representation is used in various contexts such as spinor Bose gases [8–11], entanglement classification in multiqubit systems [12–20], the Berry phase associated with the cyclic evolution of the state [21–23], investigating Lipkin-Meshkov-Glick model [24,25] and studying symmetries and properties of spin states [26–41].

The Majorana representation appears naturally in the context of $SU(2)$ coherent and anticoherent states of a given spin j [26–31]. Note that the spin coherence is not basis dependent. Properties of a coherent state $|\mathbf{n}\rangle$ of size $2j+1$ closely resemble the classical state of spin j pointing in the direction given by the vector $\mathbf{n}$. It has minimal uncertainty of spin operator $\mathbf{S}$ [42] and its Wehrl entropy [43,44] attains the minimal value [45,46]. In the Majorana representation the most coherent state is represented by one $2j$-degenerated point on a sphere. On the other hand, the most quantum, or the most anticoherent state $|\psi\rangle$ should "point nowhere" and be represented as $2j$ points equally distributed on a sphere [39–41]. However, even if the polarization (coherence) disappears, the higher moments of coherence might not vanish. Hence, anticoherence can be defined up to a given order [26]. Highly anticoherent states turned out to be applicable as optimal quantum rotosensors [32–35]. They also coincide with spherical t-designs in several dimensions [36, 38], however, a general conjecture concerning their relation was disproved [37]. Experimental realization of some of those states was discussed in [47]. The Majorana representation has proven to be a

Marcin Rudziński: marcinwojciechrudzinski@gmail.com
Adam Burchardt: adam.burchardt.uam@gmail.com
Karol Życzkowski: karol.zyczkowski@uj.edu.pl

