



Fast quantum state preparation and bath dynamics using non-Gaussian variational ansatz and quantum optimal control

Liam J. Bond,[1,2] Arghavan Safavi-Naini,[1,2] and Jiří Minář[3,1,2]

[1] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[3] *CWI, Science Park 904, 1098 XH Amsterdam, the Netherlands*

(Dated: June 5, 2023)

We combine quantum optimal control with a variational ansatz based on non-Gaussian states for fast, non-adiabatic preparation of quantum many-body states. We demonstrate this on the example of the spin-boson model, and use a multi-polaron ansatz to prepare near-critical ground states. For one mode, we achieve a reduction in infidelity of up to ≈ 60 (≈ 20) times compared to linear (optimised local adiabatic) ramps respectively; for many modes we achieve a reduction in infidelity of up to ≈ 5 times compared to non-adiabatic linear ramps. Further, we show that the typical control quantity, the leakage from the variational manifold, provides only a loose bound on the state's fidelity. Instead, in analogy to the bond dimension of matrix product states, we suggest a controlled convergence criterion based on the number of polarons. Finally, motivated by the possibility of realizations in trapped ions, we study the dynamics of a system with bath properties going beyond the paradigm of (sub/super) Ohmic couplings. We apply the ansatz to the study of the out-of-time-order-correlator (OTOC) of the bath modes in a non-perturbative regime. The scrambling time is found to be a robust feature only weakly dependent on the details of the coupling between the bath and the spin.

arXiv:2306.01730v1 [quant-ph] 2 Jun 2023

Introduction. The description of quantum systems out-of-equilibrium represents a notorious challenge. In many relevant situations one has to resort to numerical approaches ranging from non-equilibrium Monte Carlo to tensor networks [1–10]. A specific class of problems consists of systems containing bosonic degrees of freedom with an (even locally) unbounded Hilbert space. To deal with such situations, various schemes have been devised, such as path integral techniques [11–14] or effective Hamiltonian [15, 16] and lightcone conformal truncation [17–19] used predominantly in high-energy physics, which aim at describing the relevant part of the (bosonic) Hilbert space by a suitable choice of truncation procedure.

Another possibility is to exploit the continuous-variable structure of the bosonic states. Here, a novel scheme using a time-dependent variational ansatz based on *non-Gaussian* states has been recently proposed [20, 21] and successfully applied to the studies of systems ranging from Kondo impurity problem [22], central spin [23] or spin-Holstein models [24] to Bose and Fermi polarons [25–27].

In this work we demonstrate that such ansatze constitute a natural framework for the implementation of efficient state preparation schemes using quantum optimal control [28–31]. Specifically, we implement a multi-polaron ansatz [32–38] and consider the paradigmatic spin-boson model [11–14, 39–43] including in principle arbitrary couplings beyond the (sub/super) Ohmic ones and away from perturbative regimes. The choice of the spin-boson model is motivated by the fact that it plays a major role in the description of impurity problems, whilst also encompassing many platforms that are currently used for quantum simulation and computing, ranging from superconducting circuits to trapped ions [44–58]. In particular, recent realizations of the quantum Rabi-Hubbard [59] and Rabi models [60, 61] represent an ideal testbed to experimentally probe the here-presented theoretical results.

We apply the developed machinery to study (i) the onset of chaos of the bosonic bath quantified by the OTOCs, demonstrating its robustness with respect to the spin-bath couplings and (ii) fast non-adiabatic quantum many-body state preparation, including the preparation of near-critical ground states. We also highlight the limitations of the leakage as a control parameter and consider the number of polarons instead.

The model. We consider the spin-boson model, where the interaction of a two-level system with a bath of N harmonic oscillators is described by the Hamiltonian

$$H = \frac{\Delta}{2}\sigma_x + \sum_{k=1}^{N} \epsilon_k b_k^\dagger b_k - \frac{1}{2}\sigma_z \sum_{k=1}^{N} g_k (b_k^\dagger + b_k). \qquad (1)$$

Here Δ describes the tunnelling strength, ϵ_k the mode frequency and g_k the interaction between the spin and k-th mode. The operators $\sigma_{x,y,z}$ are Pauli operators acting on the spin, and b_k ($b_k^\dagger$) the annihilation (creation) operators of the bath modes satisfying $[b_k, b_{k'}^\dagger] = \delta_{kk'}$. The Hamiltonian (1) conserves the parity $P_{\text{ex}} = e^{i\pi N_{\text{ex}}}$, where $N_{\text{ex}} = 1/2(\sigma_x + 1) + \sum_k b_k^\dagger b_k$ counts the total number of excitations.

Unless stated differently, we consider Ohmic couplings, described by $\sum_k g_k^2 \delta(\omega - \epsilon_k) = 2\alpha\omega_c\omega\Theta(\omega_c - \omega)$. Here $\omega > 0$, ω_c is a high-frequency cut-off and α is a dimensionless measure of the spin-bath interaction strength. We choose a mode discretization $\epsilon_k = \omega_c k/N$, $k = 1, 2, \ldots, N$. Note that we do not enforce any restrictions