

2022–2024

arXiv:2307.11008v2 [quant-ph] 22 Nov 2024

Distillable entanglement under dually non-entangling operations

Ludovico Lami[1,2,3,*] and Bartosz Regula[4,†]

[1]*QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[2]*Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*
[3]*Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[4]*Mathematical Quantum Information RIKEN Hakubi Research Team, RIKEN Cluster for Pioneering Research (CPR) and RIKEN Center for Quantum Computing (RQC), Wako, Saitama 351-0198, Japan*

Computing the exact rate at which entanglement can be distilled from noisy quantum states is one of the longest-standing questions in quantum information. We give an exact solution for entanglement distillation under the set of dually non-entangling (DNE) operations — a relaxation of the typically considered local operations and classical communication, comprising all channels which preserve the sets of separable states and measurements. We show that the DNE distillable entanglement coincides with a modified version of the regularised relative entropy of entanglement in which the arguments are measured with a separable measurement. Ours is only the second known regularised formula for the distillable entanglement under any class of free operations in entanglement theory, after that given by Devetak and Winter for (one-way) local operations and classical communication. An immediate consequence of our finding is that, under DNE, entanglement can be distilled from any entangled state. As our second main result, we construct a general upper bound on the DNE distillable entanglement, using which we prove that the separably measured relative entropy of entanglement can be strictly smaller than the regularisation of the standard relative entropy of entanglement, solving an open problem posed by Li and Winter. Finally, we study also the reverse task of entanglement dilution and show that the restriction to DNE operations does not change the entanglement cost when compared with the larger class of non-entangling operations. This implies a strong form of irreversiblility of entanglement theory under DNE operations: even when asymptotically vanishing amounts of entanglement may be generated, entangled states cannot be converted reversibly.

INTRODUCTION

Entanglement distillation is poised to be one of the fundamental primitives of the emerging field of quantum technologies [1–3]. Its goal is to transform many copies of a noisy entangled state ρ_{AB} emitted by some source into (a smaller number of) *ebits*, i.e. perfect units of entanglement, that can then be used as fuel in a variety of quantum protocols, e.g. teleportation [4], dense coding [5], or quantum key distribution [6, 7], as well as to demonstrate violations of Bell inequalities [8]. To perform this transformation, one is allowed to use quantum operations taken from a class of *free operations* $\mathcal{F}$, assumed to comprise all quantum channels that are easy to implement on our quantum devices with the resources at our disposal. The fundamental figure of merit that characterises the ultimate efficiency of entanglement distillation on ρ_{AB} is the *distillable entanglement* under $\mathcal{F}$, denoted by $E_{d,\mathcal{F}}(\rho_{AB})$. This depends critically both on the input state ρ_{AB} and on the set of free operations $\mathcal{F}$ with which distillation is carried out.

Historically, the first class of free operations to be widely studied was that of *local operations and classical communication* (LOCC). LOCCs are the most general operations that can be performed by spatially separated parties that are bound to the rules of quantum mechanics and can communicate only classically [9]. In spite of their operational importance, LOCCs are exceedingly difficult to characterise mathematically, and as a consequence we currently do not possess any formula to compute the LOCC distillable entanglement on arbitrary states, or even decide when it is zero and when it is not [10, 11]. This is to be contrasted with the fact that we *do* possess a formula for the LOCC entanglement *cost*, which is given by the regularised entanglement of formation [12].

To bypass this problem, in recent years there has been an increasing interest in characterising entanglement manipulation under classes of free operations that approximate LOCCs. There are at least two valid reasons to pursue this goal, besides the obvious one that simpler classes of free operations offer an ideal test-bed to improve our understanding of entanglement manipulation. First, in this way one can obtain lower and upper bounds on the LOCC distillable entanglement, because if $\mathcal{F}_1 \subseteq \mathrm{LOCC} \subseteq \mathcal{F}_2$ then $E_{d,\mathcal{F}_1}(\rho_{AB}) \leq E_{d,\mathrm{LOCC}}(\rho_{AB}) \leq E_{d,\mathcal{F}_2}(\rho_{AB})$. Several of the best known bounds on the distillable entanglement can be seen as stemming from this approach. For example, the restriction to one-way LOCCs yields the hashing lower bound [13, Theorem 13], while going to the larger set of PPT operations results in the upper bounds via the negativity [14, 15] and the Rains bound [16]. Nevertheless, even under simpler classes of operations, the ultimate capabilities of entanglement manipulation — and in particular the exact value of distillable entanglement — are not known, with the exception of one-way LOCC [13].

