

2022–2025

Classifying fermionic states via many-body correlation measures

Mykola Semenyakin[1,2], Yevheniia Cheipesh[2], and Yaroslav Herasymenko[3,4,5]

[1]Perimeter Institute for Theoretical Physics, Waterloo, ON N2L 2Y5, Canada
[2]Instituut-Lorentz, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands
[3]QuSoft and CWI, Science Park 123, 1098 XG Amsterdam, The Netherlands
[4]QuTech, TU Delft, P.O. Box 5046, 2600 GA Delft, The Netherlands
[5]Delft Institute of Applied Mathematics, TU Delft, 2628 CD Delft, The Netherlands

arXiv:2309.07956v3 [quant-ph] 8 Apr 2025

Understanding the structure of quantum correlations in a many-body system is key to its computational treatment. For fermionic systems, correlations can be defined as deviations from Slater determinant states. The link between fermionic correlations and efficient computational physics methods is actively studied but remains ambiguous. We make progress in establishing this connection mathematically. In particular, we find a rigorous classification of states relative to k-fermion correlations, which admits a computational physics interpretation. Correlations are captured by a measure ω_k, a function of k-fermion reduced density matrix that we call twisted purity. A condition $\omega_k = 0$ for a given k puts the state in a class $\mathcal{G}_k$ of correlated states. Sets $\mathcal{G}_k$ are nested in k, and Slater determinants correspond to $k = 1$. Classes $\mathcal{G}_{k=\mathrm{O}(1)}$ are shown to be physically relevant, as ω_k vanishes or nearly vanishes for truncated configuration-interaction states, perturbation series around Slater determinants, and some nonperturbative eigenstates of the 1D Hubbard model. For each $k = \mathrm{O}(1)$, we give an explicit ansatz with a polynomial number of parameters that covers all states in $\mathcal{G}_k$. Potential applications of this ansatz and its connections to the coupled-cluster wavefunction are discussed.

Introduction. Quantum correlations are central to many-body quantum problems, their computational treatment, and complexity. For fermionic systems with a fixed particle number, a natural definition of uncorrelated states are Slater determinants [1]. These states arise in noninteracting systems and admit efficient computations. To characterize the correlations, or deviations of a state from a Slater determinant, quantitative measures are employed. Some measures were proposed in terms of one- and two-fermion reduced density matrices [2–6], minimal distance to the manifold of Slater states [7, 8] and so-called Slater rank [1]. Fermionic magic [9–11], measuring the deviation of a quantum circuit from fermionic linear optics [12–16], can also be considered a quantifier of correlations. Correlation measures were applied to characterize physical systems, as well as to guide computational physics and chemistry methods [3, 17–25]. Such methods are often based on approximate ansatzes, such as configuration-interaction states [30, 31], coupled-cluster ansatzes [32–36], Jastrow and Gutzwiller wavefunctions [37–39], tensor networks [27, 40–42], and generalized Gaussian ansatzes [6, 43, 44]. It is a widespread heuristic, that a state with bounded correlations should admit a representation by a compact ansatz [26–29]. The mathematical understanding of this connection, however, remains limited.

In this work, we give a classification of k-fermion correlations with a direct link to computational physics. The key new object is the twisted purity ω_k, a correlation measure which is a function of the k-body reduced density matrix, invariant under single-particle rotations. Twisted purity is also a Hermitian observable on two copies of a state. If a state yields $\omega_k = 0$, its amplitudes obey a generalization of so-called Plücker relations [45]. We denote the set of such states $\mathcal{G}_k$. Condition $\omega_{k_1} = 0$ implies $\omega_{k_2} = 0$ for all $k_2 > k_1$, so the sets $\mathcal{G}_k$ are nested in k and define a classification of fermionic states ($\mathcal{G}_1$ are Slater). We prove that each $\mathcal{G}_{k=O(1)}$ is covered by a poly-sized ansatz. This ansatz is similar to the coupled-cluster wavefunction but has a different functional form. We partially uncover the physical meaning of classes $\mathcal{G}_k$, finding that some (approximate) 1D Hubbard model eigenstates, general perturbative series around Slater determinants, and truncated configuration-interaction states belong to $\mathcal{G}_k$. By implication, twisted purity diagnoses the reducibility of a state to configuration-interaction form.

Our results complement the categorization of states via entanglement scaling [42, 46–48]. On the one hand, akin to twisted purity, entanglement can be viewed as a correlation measure, and its classification underlies computational physics methods that employ tensor networks. On the other hand, a major qualitative difference is that we focus on correlations insen-

Yaroslav Herasymenko: yaroslav@cwi.nl