

2022–2024

arXiv:2402.05793v1 [quant-ph] 8 Feb 2024

Exact quantum sensing limits for bosonic dephasing channels

Zixin Huang,[1,2,*] Ludovico Lami,[3,4,5,†] and Mark M. Wilde[6,‡]

[1] *School of Mathematical and Physical Sciences, Macquarie University, NSW 2109, Australia*
[2] *Centre for Quantum Software and Information, Faculty of Engineering and Information Technology, University of Technology Sydney, Sydney, NSW, Australia*
[3] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[4] *Korteweg–de Vries Institute for Mathematics, University of Amsterdam, Science Park 105-107, 1098 XG Amsterdam, the Netherlands*
[5] *Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*
[6] *School of Electrical and Computer Engineering, Cornell University, Ithaca, New York 14850, USA*

(Dated: February 9, 2024)

Dephasing is a prominent noise mechanism that afflicts quantum information carriers, and it is one of the main challenges towards realizing useful quantum computation, communication, and sensing. Here we consider discrimination and estimation of bosonic dephasing channels, when using the most general adaptive strategies allowed by quantum mechanics. We reduce these difficult quantum problems to simple classical ones based on the probability densities defining the bosonic dephasing channels. By doing so, we rigorously establish the optimal performance of various distinguishability and estimation tasks and construct explicit strategies to achieve this performance. To the best of our knowledge, this is the first example of a non-Gaussian bosonic channel for which there are exact solutions for these tasks.

CONTENTS

I. Introduction 1

II. Quantum channel discrimination and estimation 3
A. Quantum channel discrimination 3
B. Quantum channel estimation 3

III. Main results 4
A. Symmetric hypothesis testing 4
B. Asymmetric hypothesis testing 5
C. Quantum metrology 6

IV. Examples of bosonic dephasing channels 6

V. Optimality 6

VI. Attainability 8
A. Photon-number-superposition method 9
B. Coherent-state method 10
C. Comparison of methods for finite energy 10

VII. Bosonic dephasing and loss 11

VIII. Conclusion 11

Acknowledgments 12

References 12

A. Equivalence of hypothesis testing regions 14

B. Calculations for photon-number-superposition method 15

C. Calculations for coherent-state method 16

D. Derivation of Rician phase probability density function 17

E. Other scenarios: Strong converse exponent, error exponent, multiple channel discrimination, and antidistinguishability 18
1. Strong converse exponent 18
2. Error exponent 18
3. Multiple channel discrimination 19
4. Antidistinguishability 19

F. Multimode bosonic dephasing channels 20

G. Generalization to multiparameter channel estimation 20

H. Photon-number-superposition method in the presence of loss 20

I. INTRODUCTION

An important channel to consider in the context of quantum technologies is the bosonic dephasing channel (BDC). A single-mode BDC $\mathcal{D}_p$ is characterized by a

* zixin.huang@mq.edu.au
† ludovico.lami@gmail.com
‡ wilde@cornell.edu