



Brick Wall Quantum Circuits with Global Fermionic Symmetry

Pietro Richelli[1,2,3], Kareljan Schoutens[1,2], Alberto Zorzato[1,2*]

1 Institute for Theoretical Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

2 QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands

3 Kavli Institute of Nanoscience, Delft University of Technology, Lorentzweg 1, 2628 CJ, Delft, the Netherlands

* a.zorzato@uva.nl

February 29, 2024, revised September 9, 2024

arXiv:2402.18440v5 [quant-ph] 18 Sep 2024

Abstract

We study brick wall quantum circuits enjoying a global fermionic symmetry. The constituent 2-qubit gate, and its fermionic symmetry, derive from a 2-particle scattering matrix in integrable, supersymmetric quantum field theory in 1+1 dimensions. Our 2-qubit gate, as a function of three free parameters, is of so-called free fermionic or matchgate form, allowing us to derive the spectral structure of both the brick wall unitary $\mathbf{U_F}$ and its, non-trivial, hamiltonian limit $\mathbf{H}_\gamma$ in closed form. We find that the fermionic symmetry pins $\mathbf{H}_\gamma$ to a surface of critical points, whereas breaking that symmetry leads to non-trivial topological phases. We briefly explore quench dynamics for this class of circuits.