

arXiv:2402.18988v1 [physics.atom-ph] 29 Feb 2024

# Atom interferometry at arbitrary orientations and rotation rates

Quentin d'Armagnac de Castanet[1,2], Cyrille Des Cognets[2], Romain Arguel[2,3], Simon Templier[1], Vincent Jarlaud[1,2,*], Vincent Ménoret[1,2], Bruno Desruelle[1], Philippe Bouyer[1,4,5,6], and Baptiste Battelier[2]

[1]Exail, 1 rue François Mitterrand, 33400 Talence, France

[2]LP2N, Laboratoire Photonique, Numérique et Nanosciences, Université Bordeaux–IOGS–CNRS:UMR 5298, 1 rue François Mitterrand, 33400 Talence, France

[3]Centre National d'Etudes Spatiales, 18 avenue Edouard Belin, 31400 Toulouse, France

[4]Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam Science Park 904, 1098XH Amsterdam, Netherlands

[5]QuSoft, Science Park 123, 1098XG Amsterdam, Netherlands

[6]Eindhoven University of Technology, Eindhoven, Netherlands

[*]Corresponding author. E-mail: vincent.jarlaud@exail.com

March 1, 2024

**The exquisite precision of atom interferometers has sparked the interest of a large community for use cases ranging from fundamental physics to geodesy and inertial navigation. However, their practical use for onboard applications is still limited, not least because rotation and acceleration are intertwined in a single phase shift in free-fall atom interferometers, which makes the extraction of a useful signal more challenging. Moreover, the spatial separation of the wave packets due to rotations leads to a loss of signal. Here we present an atom interferometer operating over a large range of random angles, rotation rates and accelerations. An accurate model of the expected phase shift allows us to untangle the rotation and acceleration signals. We also implement a real-time compensation system using two fibre-optic gyroscopes and a tip-tilt platform to rotate the reference mirror and maintain the full contrast of the atom interferometer. Using these theoretical and practical tools, we reconstruct the fringes and demonstrate a single-shot sensitivity to acceleration of $24\ \mu$g, for a total interrogation time of $2T = 20\,$ms, for angles and rotation rates reaching $30°$ and $14\,°\,\mathrm{s}^{-1}$ respectively. Our hybrid rotating atom interferometer unlocks the full potential of quantum inertial sensors for onboard applications, such as autonomous navigation or gravity mapping.**