

Beating the natural Grover bound for low-energy estimation and state preparation

Harry Buhrman,[1,2,*] Sevag Gharibian,[3,†] Zeph Landau,[4,‡]
François Le Gall,[5,§] Norbert Schuch,[6,7,¶] and Suguru Tamaki[8,**]

[1] *Quantinuum, Terrington House 13-15 Hills Road Cambridge CB2 1NL, United Kingdom*
[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, The Netherlands*
[3] *Paderborn University, Department of Computer Science and Institute for Photonic Quantum Systems (PhoQS), Warburger Strasse 100, 33098 Paderborn, Germany*
[4] *University of California, Berkeley, Department of Computer Science, Berkeley, CA 94706*
[5] *Nagoya University, Graduate School of Mathematics, Furocho, Chikusaku, 464-8602 Nagoya, Japan*
[6] *University of Vienna, Faculty of Mathematics, Oskar-Morgenstern-Platz 1, 1090 Wien, Austria*
[7] *University of Vienna, Faculty of Physics, Boltzmanngasse 5, 1090 Wien, Austria*
[8] *University of Hyogo, Graduate School of Information Science, 8-2-1 Gakuennishi-machi, Nishi-ku, 651-2197 Kobe, Japan*

arXiv:2407.03073v3 [quant-ph] 22 Jul 2025

Estimating ground state energies of many-body Hamiltonians is a central task in many areas of quantum physics. In this work, we give quantum algorithms which, given any k-body Hamiltonian H, compute an estimate for the ground state energy and prepare a quantum state achieving said energy, respectively. Specifically, for any $\varepsilon > 0$, our algorithms return, with high probability, an estimate of the ground state energy of H within additive error εM, or a quantum state with the corresponding energy. Here, M is the total strength of all interaction terms, which in general is extensive in the system size. Our approach makes no assumptions about the geometry or spatial locality of interaction terms of the input Hamiltonian and thus handles even long-range or all-to-all interactions, such as in quantum chemistry, where lattice-based techniques break down. In this fully general setting, the runtime of our algorithms scales as $2^{cn/2}$ for $c < 1$, yielding the first quantum algorithms for low-energy estimation breaking a standard square root Grover speedup for unstructured search. The core of our approach is remarkably simple, and relies on showing that an extensive fraction of the interactions can be neglected with a controlled error. What this ultimately implies is that even arbitrary k-local Hamiltonians have structure in their low energy space, in the form of an exponential-dimensional low energy subspace.

Determining the properties of a quantum system at low energies is one of the central problems in the study of complex quantum many-body systems. It is most challenging in settings with long-range or all-to-all interactions, encountered for instance in the study of molecules in quantum chemistry [1, 2], but also in a variety of cold atomic and molecular as well as solid state systems [3–9]. At the same time, all-to-all interactions also appear naturally in quantum complexity theory when encoding computationally hard tasks into quantum Hamiltonians [10, 11], and would necessarily [12] have to be present in the sought-for quantum PCP theorem [13].

The ability of both classical and quantum algorithms to compute low-energy properties of quantum many-body systems is fundamentally limited by quantum complexity theory, which poses restrictions on what generally applicable algorithms can provably achieve. Finding the ground state energy of a system of n spins up to precision $1/\mathrm{poly}(n)$ is known to be QMA-hard (where QMA is the quantum version of NP) [10, 11, 14–16], and thus hard even for quantum computers.

To make the problem more tractable, one can aim for some given *extensive accuracy* in terms of the total energy scale. If the total strength of the interactions scales like the system size, as expected in physical systems, this amounts to a constant (but arbitrarily small) energy density, and thus allows to study properties of the system which are stable at finite temperature. For systems with spatially local interactions, e.g., on a lattice, this problem can be solved efficiently (for a fixed precision) by cutting the system into constant-sized patches and neglecting the couplings between those patches [12]. However, this is no longer possible for sytems with all-to-all interactions. In fact, the problem must remain at least NP-hard, as the PCP theorem implies that constraint satisfaction problems (which map to Hamiltonians with long-range couplings) remain NP-hard even for determining whether a fixed fraction of terms is violated [17, 18]; the validity of a potential quantum PCP theorem [13] would even imply its QMA-hardness.

In the light of these hardness results and the unstructured nature of the problem, the best runtime we can expect for a classical algorithm for estimating properties at constant energy density is exponential, by exact diagonalization methods such as the Lanczos algorithm [19, 20]. In fact, the Strong Exponential Time Hypothesis states that classical algorithms *must* run essentially in time scaling with 2^n [21]. On a quantum computer, we can use Grover search, leading to a quadratic speedup and thus a runtime of $O(2^{n/2})$ for qubit systems [22–25]. These methods can be improved if we are given access to a *trial* or *guiding* state for the low-energy subspace, as in recent quantum algorithms for ground state energy estimation [26, 27]; however, the computation of such guiding states is a serious bottleneck, e.g., in current quantum chemistry applications [28].