



Vibrationally coupled Rydberg atom-ion molecules

Ilango Maran,[1] Liam J. Bond,[2,3] Jeremy T. Young,[3] Arghavan Safavi-Naini,[2,3] and Rene Gerritsma[1,2]

[1] *Van der Waals-Zeeman Institute, Institute of Physics, University of Amsterdam, 1098 XH Amsterdam, the Netherlands*
[2] *QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands*
[3] *Institute for Theoretical Physics, Institute of Physics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, the Netherlands*

(Dated: May 13, 2025)

We study the occurrence of Rydberg atom-ion molecules (RAIMs) in a hybrid atom-ion system with an ion crystal trapped in a Paul trap coupled to Rydberg atoms on its either ends. To assess the feasibility of such a system, we perform a detailed Floquet analysis of the effect of the Paul trap's rf potential on the RAIMs and provide a qualitative analysis of the survival probability based on scaling laws. We conclude that the RAIM survives for sufficiently weak and low frequency traps. We then use this hybrid system and propose a scheme to utilise the common motional modes of the ion crystal to suppress (blockade) or enhance (anti-blockade) the probability of forming two RAIMs at the ends of the chain, replacing the typical blockade radius by the length of the ion crystal.

arXiv:2409.13469v2 [quant-ph] 12 May 2025

Recently, a new type of Rydberg molecule consisting of an ion and a Rydberg atom was proposed and observed in an ultracold cloud of rubidium atoms [1–4]. The binding mechanism is based on the flipping dipole induced in the Rydberg atom by the ion's electric field, which results in an atom-ion separation-dependent mixing of the different Rydberg states [5–12]. Molecular bound states are found in the potential wells located at the avoided crossings between the low orbital angular momentum Rydberg state (nS and nP) and the hydrogenic manifolds. These molecular ions can have remarkably large bond lengths on the order of a few μm and binding energies as low as a few MHz [1–4].

A natural question is whether we can extend the Rydberg atom-ion molecule to include more particles. In this letter, we study the possibility of replacing the single ion with a trapped ion crystal. We show that the Rydberg atom-ion molecule (RAIM) survives the effects of the Paul trap, provided the trap is sufficiently weak and low-frequency, and utilize the rich vibrational spectrum added by the additional ions to propose novel longer-range Rydberg blockade and anti-blockade mechanisms.

The structure of the paper is as follows. We illustrate the system in Fig. 1(a)(i), where a two ion chain interacts with Rydberg atoms placed at either end, to form two RAIMs. In Fig. 1(b), we plot the potential energy curves due to the Coulombic interaction between the ion and the Rydberg atom, assuming that the atomic core and the ion are stationary compared to the Rydberg electron [13]. The potential wells support RAIMs at specific values of the core-ion separation r_{ci}, one to three orders of magnitude smaller than the typical ion-ion separation in the crystal. Next, we use Floquet formalism to study the effect of the Paul trap's rf-drive on the RAIM, which can potentially induce transitions from the molecular state to unbound atomic states, as shown in Fig. 1(c). We use a simple qualitative model using scaling laws to understand the RAIM's survival probability and identify a broad range of Rydberg states and Paul trap drive frequencies where trap-induced losses will be negligible. Finally, we use the common motional modes of the trapped ion crystal to mediate interactions between the two Ry-




FIG. 1. (a) Two vibrationally coupled RAIMs aligned along the rf-null axis of a Paul trap potential in the xy-plane (axial case) where z_1, z_2, ζ_1 and ζ_2 represent the positions of the ions and the cores of the Rydberg atoms respectively is shown in (i). A RAIM aligned along the z-axis of a Paul trap potential in the zy-plane (radial case) is shown in (ii). (b) Potential energy curves of a $^{87}\mathrm{Rb}^{9}\mathrm{Be}^{+}$ RAIM for $|m_j| = 1/2$ in the vicinity of the $22P$ state is shown above and a magnified view of a part of the $22P_{1/2}$ potential well with its first five bound states is shown below. The energies are relative to the field-free atomic $22P_{3/2}$ state. (c) Schematic depicting a Rydberg atom escaping the well state by coupling with another state through a high-order photon process. (d) Energy levels of the four possible electronic states of the two atoms, each with its own ladder of motional states characterized by their respective phonon occupation numbers that can be exploited to mediate long-range Rydberg interactions.