



2021–2025

Classical versus quantum queries in quantum PCPs with classical proofs

Harry Buhrman[1,2], François Le Gall[3], and Jordi Weggemans[4]

[1]Quantinuum, United Kingdom
[2]QuSoft & University of Amsterdam, Amsterdam, The Netherlands
[3]Nagoya University, Japan
[4]QuSoft & CWI, Amsterdam, the Netherlands

November 5, 2024

arXiv:2411.00946v1 [quant-ph] 1 Nov 2024

Abstract

We generalize quantum-classical PCPs, first introduced by Weggemans, Folkertsma and Cade (TQC 2024), to allow for q *quantum* queries to a polynomially-sized classical proof ($\mathsf{QCPCP}_{Q,c,s}[q]$). Exploiting a connection with the polynomial method, we prove that for any constant q, promise gap $c - s = \Omega(1/\mathrm{poly}(n))$ and $\delta > 0$, we have $\mathsf{QCPCP}_{Q,c,s}[q] \subseteq \mathsf{QCPCP}_{1-\delta,1/2+\delta}[3] \subseteq \mathsf{BQ}\cdot\mathsf{NP}$, where $\mathsf{BQ}\cdot\mathsf{NP}$ is the class of promise problems with quantum reductions to an NP-complete problem. Surprisingly, this shows that we can amplify the promise gap from inverse polynomial to constant for constant query quantum-classical PCPs, and that any quantum-classical PCP making any constant number of quantum queries can be simulated by one that makes only three classical queries. Nevertheless, even though we can achieve promise gap amplification, our result also gives strong evidence that there exists no constant query quantum-classical PCP for QCMA, as it is unlikely that $\mathsf{QCMA} \subseteq \mathsf{BQ}\cdot\mathsf{NP}$, which we support by giving oracular evidence. In the (poly-)logarithmic query regime, we show for any positive integer c, there exists an oracle relative to which $\mathsf{QCPCP}[\mathcal{O}(\log^c n)] \subsetneq \mathsf{QCPCP}_Q[\mathcal{O}(\log^c n)]$, contrasting the constant query case where the equivalence of both query models holds relative to any oracle. Finally, we connect our results to more general quantum-classical interactive proof systems.

Contents

1 Introduction **2**
1.1 Our results . . . 2
1.2 Proof ideas . . . 4
1.3 Implication to quantum-classical interactive proof systems . . . 5
1.4 Discussion and open questions . . . 6

2 Preliminaries **6**
2.1 Complexity theory . . . 6
2.2 The OR ∘ Forrelation oracle . . . 7

3 Quantum reductions and the BQ-operator **8**
3.1 Randomized reductions and Schöning's BP-operator . . . 9
3.2 The BQ-operator . . . 10

4 Pulling the quantumness out of quantum-classical PCPs **12**
4.1 A threshold problem for multi-linear polynomials . . . 13
4.2 The polynomial method . . . 14
4.3 The quantum reduction . . . 15
4.4 Implications . . . 20

5 Oracle separations for quantum-classical PCPs **24**
5.1 The quantum-classical PCP conjecture is false relative to an oracle . . . 24
5.2 Oracle separations for logarithmic queries . . . 25

A Omitted proofs **29**

